



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TRAM2 siRNA (m): sc-154580

BACKGROUND

TRAM2 (translocation associated membrane protein 2) is a 370 amino acid multi-pass membrane protein containing one TLC (TRAM/LAG1/CLN8) domain. TRAM2 interacts with SERCA2B and COL1A1 and is necessary for collagen type I synthesis. TRAM2 facilitates proper folding of collagen by coupling the activity of SERCA2B, an ER Ca²⁺ pump, and the activity of translocons, thereby increasing local Ca²⁺ concentration at the site of collagen synthesis and stimulating molecular chaperones that are involved in collagen folding. It is suggested that TRAM2 is regulated by RUNX2 in a BMP-dependent manner and may play a role in the overall osteogenic function of RUNX2. RUNX2 is essential for skeletal mineralization in which it stimulates osteoblast differentiation of mesenchymal stem cells, promotes chondrocyte hypertrophy and contributes to endothelial cell migration and vascular invasion of developing bones.

REFERENCES

1. Onuchic, L.F., et al. 1999. Genomic organization of the KIAA0057 gene that encodes a TRAM-like protein and its exclusion as a polycystic kidney and hepatic disease 1 (PKHD1) candidate gene. *Mamm. Genome* 10: 1175-1178.
2. Stefanovic, B., et al. 2004. TRAM2 protein interacts with endoplasmic reticulum Ca²⁺ pump Serca2b and is necessary for collagen type I synthesis. *Mol. Cell. Biol.* 24: 1758-1768.
4. Yoshida, C.A., et al. 2005. Role of Runx proteins in chondrogenesis. *Crit. Rev. Eukaryot. Gene Expr.* 15: 243-254.
5. Westendorf, J.J. 2006. Transcriptional co-repressors of Runx2. *J. Cell. Biochem.* 98: 54-64.
6. Pregizer, S., et al. 2007. Identification of novel Runx2 targets in osteoblasts: cell type-specific BMP-dependent regulation of Tram2. *J. Cell. Biochem.* 102: 1458-1471.
7. Poot, M., et al. 2009. Three *de novo* losses and one insertion within a pericentric inversion of chromosome 6 in a patient with complete absence of expressive speech and reduced pain perception. *Eur. J. Med. Genet.* 52: 27-30.

CHROMOSOMAL LOCATION

Genetic locus: Tram2 (mouse) mapping to 1 A4.

PRODUCT

TRAM2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRAM2 shRNA Plasmid (m): sc-154580-SH and TRAM2 shRNA (m) Lentiviral Particles: sc-154580-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRAM2 siRNA (m) is recommended for the inhibition of TRAM2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRAM2 gene expression knockdown using RT-PCR Primer: TRAM2 (m)-PR: sc-154580-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.