



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TRUB2 siRNA (m): sc-154700

BACKGROUND

TRUB2 (tRNA pseudouridine synthase 2) is a 331 amino acid protein that most likely is responsible for the synthesis of pseudouridine from isomerization of uracil in tRNA. TRUB2 contains one TruB domain and shares 30% sequence similarity with TRUB1. Both TRUB1 and TRUB2 are phylogenetically linked to the original eubacterial tRNA synthases and distinct from Dyskerin. The gene encoding TRUB2 maps to human chromosome 9q34.11, which houses over 900 genes and comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

1. Nurse, K., Wrzesinski, J., Bakin, A., Lane, B.G. and Ofengand, J. 1995. Purification, cloning, and properties of the tRNA psi 55 synthase from *Escherichia coli*. *RNA* 1: 102-112.
2. Kinghorn, S.M., O'Byrne, C.P., Booth, I.R. and Stansfield, I. 2002. Physiological analysis of the role of truB in *Escherichia coli*: a role for tRNA modification in extreme temperature resistance. *Microbiology* 148: 3511-3520.
3. Zucchini, C., Strippoli, P., Biolchi, A., Solmi, R., Lenzi, L., D'Addabbo, P., Carinci, P. and Valvassori, L. 2003. The human TruB family of pseudouridine synthase genes, including the dyskeratosis congenita 1 gene and the novel member TRUB1. *Int. J. Mol. Med.* 11: 697-704.
4. Pan, H., Agarwalla, S., Moustakas, D.T., Finer-Moore, J. and Stroud, R.M. 2003. Structure of tRNA pseudouridine synthase TruB and its RNA complex: RNA recognition through a combination of rigid docking and induced fit. *Proc. Natl. Acad. Sci. USA* 100: 12648-12653.
5. Roovers, M., Hale, C., Tricot, C., Terns, M.P., Terns, R.M., Grosjean, H. and Droogmans, L. 2006. Formation of the conserved pseudouridine at position 55 in archaeal tRNA. *Nucleic Acids Res.* 34: 4293-4301.
6. Gurha, P., Joardar, A., Chaurasia, P. and Gupta, R. 2007. Differential roles of archaeal box H/ACA proteins in guide RNA-dependent and independent pseudouridine formation. *RNA Biol.* 4: 101-109.
7. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610727. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Trub2 (mouse) mapping to 2 B.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TRUB2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRUB2 shRNA Plasmid (m): sc-154700-SH and TRUB2 shRNA (m) Lentiviral Particles: sc-154700-V as alternate gene silencing products.

For independent verification of TRUB2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154700A, sc-154700B and sc-154700C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRUB2 siRNA (m) is recommended for the inhibition of TRUB2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRUB2 gene expression knockdown using RT-PCR Primer: TRUB2 (m)-PR: sc-154700-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.