



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# TSPAN11 siRNA (m): sc-154724

## BACKGROUND

Tetraspanins are a group of hydrophobic membrane proteins that interact with a wide variety of proteins including intracellular signaling molecules, integrins and membrane receptors. Members of the tetraspanin family are characterized by the presence of four hydrophobic domains and play a role in cell development, activation, growth and motility. TSPAN11 (tetraspanin-11) is a 253 amino acid multi-pass membrane protein belonging to the tetraspanin (TM4SF) family. The gene encoding TSPAN11 maps to human chromosome 12p11.21, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

## REFERENCES

1. Todd, S.C., Doctor, V.S. and Levy, S. 1998. Sequences and expression of six new members of the tetraspanin/TM4SF family. *Biochim. Biophys. Acta* 1399: 101-104.
2. Domínguez-Jimenez, C., Yáñez-Mó, M., Carreira, A., Tejedor, R., González-Amaro, R., Alvarez, V. and Sánchez-Madrid, F. 2001. Involvement of  $\alpha 3$  integrin/tetraspanin complexes in the angiogenic response induced by angiotensin II. *FASEB J.* 15: 1457-1459.
3. Berditchevski, F. 2001. Complexes of tetraspanins with integrins: more than meets the eye. *J. Cell Sci.* 114: 4143-4151.
4. Yokoyama, T., Nakatani, S. and Murakami, A. 2003. A case of Kniest dysplasia with retinal detachment and the mutation analysis. *Am. J. Ophthalmol.* 136: 1186-1188.
5. Chen, L., Li, X., Wang, G.L., Wang, Y., Zhu, Y.Y. and Zhu, J. 2008. Clinicopathological significance of overexpression of TSPAN1, Ki67 and CD34 in gastric carcinoma. *Tumori* 94: 531-538.
6. Benussi, D.G., Costa, P., Zollino, M., Murdolo, M., Petix, V., Carrozza, M. and Pecile, V. 2009. Trisomy 12p and monosomy 4p: phenotype-genotype correlation. *Genet. Test. Mol. Biomarkers* 13: 199-204.
7. Scholz, C.J., Kurzeder, C., Koretz, K., Windisch, J., Kreienberg, R., Sauer, G. and Deissler, H. 2009. Tspan-1 is a tetraspanin preferentially expressed by mucinous and endometrioid subtypes of human ovarian carcinomas. *Cancer Lett.* 275: 198-203.
8. Chen, L., Zhu, Y.Y., Zhang, X.J., Wang, G.L., Li, X.Y., He, S., Zhang, J.B. and Zhu, J.W. 2009. TSPAN1 protein expression: a significant prognostic indicator for patients with colorectal adenocarcinoma. *World J. Gastroenterol.* 15: 2270-2276.
9. Bennett, G., Sadlier, D., Doran, P.P., Macmathuna, P. and Murray, D.W. 2011. A functional and transcriptomic analysis of NET1 bioactivity in gastric cancer. *BMC Cancer* 11: 50.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Tspan11 (mouse) mapping to 6 F3.

## PRODUCT

TSPAN11 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TSPAN11 shRNA Plasmid (m): sc-154724-SH and TSPAN11 shRNA (m) Lentiviral Particles: sc-154724-V as alternate gene silencing products.

For independent verification of TSPAN11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154724A, sc-154724B and sc-154724C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TSPAN11 siRNA (m) is recommended for the inhibition of TSPAN11 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TSPAN11 gene expression knockdown using RT-PCR Primer: TSPAN11 (m)-PR: sc-154724-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60°C and the extension temperature should be 68-72°C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.