



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# UBA52 siRNA (m): sc-154840

## BACKGROUND

Ubiquitin (Ub) is among the most phylogenetically conserved proteins known. The primary function of ubiquitin is to clear abnormal, foreign and improperly folded proteins by targeting them for degradation by the 26S proteasome. Encoded by four genes, ubiquitin is synthesized as precursor proteins that consist of either single ubiquitin moieties fused 5' to unrelated carboxyl extension proteins, known as UBA type, or polyubiquitin chains that are cleaved into moieties of the UBB or UBC types. As a UBA type ubiquitin, UBA52 (Ubiquitin-60S ribosomal protein L40), also known as CEP52 and Ubiquitin A-52 residue ribosomal protein fusion product 1, is a 128 amino acid protein that is cleaved into ubiquitin and 60S ribosomal protein L40. Normally expressed in lymphocytes and placenta, UBA52 is overexpressed in renal cancer cells and colorectal carcinoma, suggesting it may have a role to play in tumorigenesis.

## REFERENCES

1. Salvesen, G., Lloyd, C. and Farley, D. 1987. cDNA encoding a human homolog of yeast ubiquitin 1. *Nucleic Acids Res.* 15: 5485.
2. Kanayama, H., Tanaka, K., Aki, M., Kagawa, S., Miyaji, H., Satoh, M., Okada, F., Sato, S., Shimbara, N. and Ichihara, A. 1991. Changes in expressions of proteasome and ubiquitin genes in human renal cancer cells. *Cancer Res.* 51: 6677-6685.
3. Baker, R.T. and Board, P.G. 1991. The human ubiquitin-52 amino acid fusion protein gene shares several structural features with mammalian ribosomal protein genes. *Nucleic Acids Res.* 19: 1035-1040.
4. Barnard, G.F., Mori, M., Staniunas, R.J., Begum, N.A., Bao, S., Puder, M., Cobb, J., Redman, K.L., Steele, G.D. and Chen, L.B. 1995. Ubiquitin fusion proteins are overexpressed in colon cancer but not in gastric cancer. *Biochim. Biophys. Acta* 1272: 147-153.
5. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 191321. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Okumura, F., Hatakeyama, S., Matsumoto, M., Kamura, T. and Nakayama, K.I. 2004. Functional regulation of FEZ1 by the U-box-type ubiquitin ligase E4B contributes to neuritogenesis. *J. Biol. Chem.* 279: 53533-53543.
7. Cripps, D., Thomas, S.N., Jeng, Y., Yang, F., Davies, P. and Yang, A.J. 2006. Alzheimer disease-specific conformation of hyperphosphorylated paired helical filament-Tau is polyubiquitinated through Lys-48, Lys-11, and Lys-6 ubiquitin conjugation. *J. Biol. Chem.* 281: 10825-10838.
8. Huang, F., Kirkpatrick, D., Jiang, X., Gygi, S. and Sorkin, A. 2006. Differential regulation of EGF receptor internalization and degradation by multiubiquitination within the kinase domain. *Mol. Cell* 21: 737-748.
9. Motegi, A., Liaw, H.J., Lee, K.Y., Roest, H.P., Maas, A., Wu, X., Moinova, H., Markowitz, S.D., Ding, H., Hoeijmakers, J.H. and Myung, K. 2008. Polyubiquitination of proliferating cell nuclear antigen by HILTF and SHPRH prevents genomic instability from stalled replication forks. *Proc. Natl. Acad. Sci. USA* 105: 12411-12416.

## CHROMOSOMAL LOCATION

Genetic locus: Uba52 (mouse) mapping to 8 B3.3.

## PRODUCT

UBA52 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UBA52 shRNA Plasmid (m): sc-154840-SH and UBA52 shRNA (m) Lentiviral Particles: sc-154840-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

UBA52 siRNA (m) is recommended for the inhibition of UBA52 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor UBA52 gene expression knockdown using RT-PCR Primer: UBA52 (m)-PR: sc-154840-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.