



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



## XKR4 siRNA (m): sc-155373

### BACKGROUND

Kell and XK are two covalently linked plasma membrane proteins that constitute the Kell blood group system, a group of antigens on the surface of red blood cells that are important determinants of blood type and targets for autoimmune or alloimmune diseases. XK is a 444 amino acid protein that spans the membrane ten times and carries the ubiquitous antigen, Kx, which determines blood type. The XK (X-linked Kx blood group)-related gene family are homologs of XK. XKR4 (XK-related protein 4) is a 650 amino acid multi-pass membrane protein that likely is a component of the XK/Kell complex of the Kell blood group system. The gene encoding XKR4 maps to human chromosome 8, which is made up of nearly 146 million bases and encodes about 800 genes. Single nucleotide polymorphisms (SNPs) of the XKR4 gene are associated with iloperidone efficacy, a antipsychotic drug that is used to treat schizophrenic patients.

### REFERENCES

1. Lee, S., et al. 2000. Functional and structural aspects of the Kell blood group system. *Transfus. Med. Rev.* 14: 93-103.
2. Lee, S., et al. 2000. The Kell blood group system: Kell and XK membrane proteins. *Semin. Hematol.* 37: 113-121.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 314850. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Singleton, B.K., et al. 2003. McLeod syndrome resulting from a novel XK mutation. *Br. J. Haematol.* 122: 682-685.
5. Feder, M., et al. 2005. Identification of a new family of putative PD-(D/E)XK nucleases with unusual phylogenomic distribution and a new type of the active site. *BMC Genomics* 6: 21.
6. Pu, J.J., et al. 2005. Onset of expression of the components of the Kell blood group complex. *Transfusion* 45: 969-974.
7. Starling, A., et al. 2005. A family with McLeod syndrome and calpainopathy with clinically overlapping diseases. *Neurology* 65: 1832-1833.
8. Zeman, A., et al. 2005. McLeod syndrome: life-long neuropsychiatric disorder due to a novel mutation of the XK gene. *Psychiatr. Genet.* 15: 291-293.
9. Lavedan, C., et al. 2009. Association of the NPAS3 gene and five other loci with response to the antipsychotic iloperidone identified in a whole genome association study. *Mol. Psychiatry* 14: 804-819.

### CHROMOSOMAL LOCATION

Genetic locus: Xkr4 (mouse) mapping to 1 A1.

### PRODUCT

XKR4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see XKR4 shRNA Plasmid (m): sc-155373-SH and XKR4 shRNA (m) Lentiviral Particles: sc-155373-V as alternate gene silencing products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

XKR4 siRNA (m) is recommended for the inhibition of XKR4 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor XKR4 gene expression knockdown using RT-PCR Primer: XKR4 (m)-PR: sc-155373-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.