



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# ZFY1 siRNA (m): sc-155600



The Power to Question

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZFY1 (zinc finger protein 1, Y linked), also known ZFY1 is a 782 amino acid nuclear protein belonging to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family and the ZFX/ZFY subfamily. Containing 13 C<sub>2</sub>H<sub>2</sub>-type zinc fingers, ZFY1 is expressed in the genital ridge and adult testis and may be a probable transcription activator. The gene encoding ZFY1 maps to mouse chromosome Y A1.

## REFERENCES

- Ashworth, A., Swift, S. and Affara, N. 1989. Sequence of cDNA for murine Zfy-1, a candidate for Tdy. *Nucleic Acids Res.* 17: 2864.
- Hayashida, H., Kuma, K. and Miyata, T. 1992. Interchromosomal gene conversion as a possible mechanism for explaining divergence patterns of ZFY-related genes. *J. Mol. Evol.* 35: 181-183.
- Pamilo, P. and Bianchi, N.O. 1993. Evolution of the Zfx and Zfy genes: rates and interdependence between the genes. *Mol. Biol. Evol.* 10: 271-281.
- Zambrowicz, B.P., Zimmermann, J.W., Harendza, C.J., Simpson, E.M., Page, D.C., Brinster, R.L. and Palmiter, R.D. 1994. Expression of a mouse Zfy-1/lacZ transgene in the somatic cells of the embryonic gonad and germ cells of the adult testis. *Development* 120: 1549-1559.
- Zambrowicz, B.P., Findley, S.D., Simpson, E.M., Page, D.C. and Palmiter, R.D. 1994. Characterization of the murine Zfy1 and Zfy2 promoters. *Genomics* 24: 406-408.
- Mahaffey, C.L., Bayleran, J.K., Yeh, G.Y., Lee, T.C., Page, D.C. and Simpson, E.M. 1997. Intron/exon structure confirms that mouse Zfy1 and Zfy2 are members of the ZFY gene family. *Genomics* 41: 123-127.
- Nagamine, C.M. and Carlisle, C. 1997. Zfy2/1 fusion gene fails to replicate Zfy1 expression pattern in fetal gonads. *Genomics* 43: 397-398.
- Johnston, C.M., Shimeld, S.M. and Sharpe, P.T. 1998. Molecular evolution of the ZFY and ZNF6 gene families. *Mol. Biol. Evol.* 15: 129-137.
- Hansen, M.A., Nielsen, J.E., Tanaka, M., Almstrup, K., Skakkebaek, N.E. and Leffers, H. 2006. Identification and expression profiling of 10 novel spermatid expressed CYPT genes. *Mol. Reprod. Dev.* 73: 568-579.

## CHROMOSOMAL LOCATION

Genetic locus: Zfy1 (mouse) mapping to Y A1.

## PRODUCT

ZFY1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZFY1 shRNA Plasmid (m): sc-155600-SH and ZFY1 shRNA (m) Lentiviral Particles: sc-155600-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZFY1 siRNA (m) is recommended for the inhibition of ZFY1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZFY1 gene expression knockdown using RT-PCR Primer: ZFY1 (m)-PR: sc-155600-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.