



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ZNF426 siRNA (m): sc-155715



The Power to Question

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF426 (zinc finger protein 426), also known as MGC2663, is a 554 amino acid protein that is thought to be involved in transcriptional regulation. Localized to the nucleus, ZNF426 contains one KRAB domain and 12 C₂H₂-type zinc fingers through which it may convey DNA, RNA and protein binding capabilities. Specifically, ZNF426 may interact with the viral protein KSHV ORF 50 and, through this interaction, may activate viral gene transcription.

REFERENCES

1. Huebner, K., Druck, T., Croce, C.M. and Thiesen, H.J. 1991. Twenty-seven nonoverlapping zinc finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726-740.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Rousseau-Merck, M.F., Hillion, J., Jonveaux, P., Couillin, P., Sete, P., Thiesen, H.J. and Berger, R. 1994. Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583-587.
4. Williams, A.J., Blacklow, S.C. and Collins, T. 1999. The zinc finger-associated SCAN box is a conserved oligomerization domain. *Mol. Cell. Biol.* 19: 8526-8535.
5. Wang, S., Liu, S., Wu, M.H., Geng, Y. and Wood, C. 2001. Identification of a cellular protein that interacts and synergizes with the RTA (ORF50) protein of Kaposi's sarcoma-associated herpesvirus in transcriptional activation. *J. Virol.* 75: 11961-11973.
6. Shannon, M., Hamilton, A.T., Gordon, L., Branscomb, E. and Stubbs, L. 2003. Differential expansion of zinc-finger transcription factor loci in homologous human and mouse gene clusters. *Genome Res.* 13: 1097-1110.
7. Englbrecht, C.C., Schoof, H. and Böhm, S. 2004. Conservation, diversification and expansion of C₂H₂ zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39.
8. Yang, Z. and Wood, C. 2007. The transcriptional repressor K-RBP modulates RTA-mediated transactivation and lytic replication of Kaposi's sarcoma-associated herpesvirus. *J. Virol.* 81: 6294-6306.
9. Yang, Z., Yan, Z. and Wood, C. 2008. Kaposi's sarcoma-associated herpesvirus transactivator RTA promotes degradation of the repressors to regulate viral lytic replication. *J. Virol.* 82: 3590-3603.

CHROMOSOMAL LOCATION

Genetic locus: Zfp426 (mouse) mapping to 9 A3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

ZNF426 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF426 shRNA Plasmid (m): sc-155715-SH and ZNF426 shRNA (m) Lentiviral Particles: sc-155715-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF426 siRNA (m) is recommended for the inhibition of ZNF426 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF426 gene expression knockdown using RT-PCR Primer: ZNF426 (m)-PR: sc-155715-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.