



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# ZNF7 siRNA (m): sc-155782



The Power to Question

## BACKGROUND

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF7 (zinc finger protein 7), also known as KOX4 or HF.16, is a 686 amino acid zinc-finger protein that belongs to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc finger family. Localized to the nucleus, ZFP3 contains 15 C<sub>2</sub>H<sub>2</sub>-type zinc fingers and is thought to play a role in transcriptional regulation.

## REFERENCES

1. Lania, L., Donti, E., Pannuti, A., Pascucci, A., Pengue, G., Feliciello, I., La Mantia, G., Lanfrancone, L. and Pelicci, P.G. 1990. cDNA isolation, expression analysis, and chromosomal localization of two human zinc-finger genes. *Genomics* 6: 333-340.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Bray, P., Lichter, P., Thiesen, H.J., Ward, D.C. and Dawid, I.B. 1991. Characterization and mapping of human genes encoding zinc-finger proteins. *Proc. Natl. Acad. Sci. USA* 88: 9563-9567.
4. Huebner, K., Druck, T., LaForgia, S., Lasota, J., Croce, C.M., Lanfrancone, L., Donti, E., Pengue, G., La Mantia, G. and Pelicci, P.G. 1993. Chromosomal localization of four human zinc finger cDNAs. *Hum. Genet.* 91: 217-222.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 194531. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: Zfp7 (mouse) mapping to 15 D3.

## PRODUCT

ZNF7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF7 shRNA Plasmid (m): sc-155782-SH and ZNF7 shRNA (m) Lentiviral Particles: sc-155782-V as alternate gene silencing products.

For independent verification of ZNF7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155782A, sc-155782B and sc-155782C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZNF7 siRNA (m) is recommended for the inhibition of ZNF7 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF7 gene expression knockdown using RT-PCR Primer: ZNF7 (m)-PR: sc-155782-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.