



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# ZNF711 siRNA (m): sc-155786



The Power to Question

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF6, also known as ZNF711 (zinc finger protein 711), MRX97 or CMPX1, is a 761 amino acid protein that contains 12 C<sub>2</sub>H<sub>2</sub>-type zinc fingers and belongs to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family. Localizing to the nucleus, ZNF6 is expressed in neural tissues and becomes phosphorylated upon DNA damage. Acting as a transcription regulator required for proper brain development, defects to ZNF6 have been linked to mental retardation X-linked ZNF711-related (MRXZ). Existing as three alternatively spliced isoforms, the gene encoding ZNF6 maps to human chromosome Xq21.1.

## REFERENCES

- Lloyd, S.L., Sargent, C.A., Chalmers, J., Lim, E., Habeebu, S.S. and Affara, N.A. 1991. An X-linked zinc finger gene mapping to Xq21.1-q21.3 closely related to ZFX and ZFY: possible origins from a common ancestral gene. *Nucleic Acids Res.* 19: 4835-4841.
- Colleaux, L., May, M., Belougne, J., Lepaslier, D., Schwartz, C. and Fontes, M. 1996. Localisation of two candidate genes for mental retardation using a YAC physical map of the Xq21.1-21.2 subbands. *J. Med. Genet.* 33: 353-357.
- Johnston, C.M., Shimeld, S.M. and Sharpe, P.T. 1998. Molecular evolution of the ZFY and ZNF6 gene families. *Mol. Biol. Evol.* 15: 129-137.
- Matsuoka, S., Ballif, B.A., Smogorzewska, A., McDonald, E.R., Hurov, K.E., Luo, J., Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y., Shiloh, Y., Gygi, S.P. and Elledge, S.J. 2007. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316: 1160-1166.
- Tarpey, P.S., Smith, R., Pleasance, E., Whibley, A., Edkins, S., Hardy, C., O'Meara, S., Latimer, C., Dicks, E., Menzies, A., Stephens, P., Blow, M., Greenman, C., Xue, Y., Tyler-Smith, C., Thompson, D., Gray, K., et al. 2009. A systematic, large-scale resequencing screen of X-chromosome coding exons in mental retardation. *Nat. Genet.* 41: 535-543.
- Kleine-Kohlbrecher, D., Christensen, J., Vandamme, J., Abarrategui, I., Bak, M., Tommerup, N., Shi, X., Gozani, O., Rappaport, J., Salcini, A.E. and Helin, K. 2010. A functional link between the histone demethylase PHF8 and the transcription factor ZNF711 in X-linked mental retardation. *Mol. Cell* 38: 165-178.
- Kantajärvi, K., Kotala, I., Rehnström, K., Ylisaukko-Oja, T., Vanhala, R., von Wendt, T.N., von Wendt, L. and Järvelä, I. 2011. Fine mapping of Xq11.1-q21.33 and mutation screening of RPS6KA6, ZNF711, ACSL4, DLG3, and IL1RAPL2 for autism spectrum disorders (ASD). *Autism Res.* 4: 228-233.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Zfp711 (mouse) mapping to X E1.

## PRODUCT

ZNF711 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF711 shRNA Plasmid (m): sc-155786-SH and ZNF711 shRNA (m) Lentiviral Particles: sc-155786-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZNF711 siRNA (m) is recommended for the inhibition of ZNF711 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF711 gene expression knockdown using RT-PCR Primer: ZNF711 (m)-PR: sc-155786-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.