



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ZNF879 siRNA (h): sc-155996



The Power to Question

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF879 (zinc finger protein 879) is a 563 amino acid protein belonging to the Krüppel C₂H₂-type zinc-finger protein family. ZNF879 contains 13 C₂H₂-type zinc fingers and one KRAB domain. This nuclear protein is likely involved in transcriptional regulation. The gene encoding ZNF879 maps to human chromosome 5q35.3, which contains 181 million base pairs and comprises nearly 6% of the human genome. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

1. Freemont, P.S. 1993. The RING finger. A novel protein sequence motif related to the zinc finger. *Ann. N.Y. Acad. Sci.* 684: 174-192.
2. Klug, A. 1999. Zinc finger peptides for the regulation of gene expression. *J. Mol. Biol.* 293: 215-218.
3. Schafer, I.A., Robin, N.H., Posch, J.J., Clark, B.A., Izumo, S. and Schwartz, S. 2001. Distal 5q deletion syndrome: phenotypic correlations. *Am. J. Med. Genet.* 103: 63-68.
4. Laity, J.H., Lee, B.M. and Wright, P.E. 2001. Zinc finger proteins: new insights into structural and functional diversity. *Curr. Opin. Struct. Biol.* 11: 39-46.
5. Mainardi, P.C., Perfumo, C., Cali, A., Coucourde, G., Pastore, G., Cavani, S., Zara, F., Overhauser, J., Pierluigi, M. and Bricarelli, F.D. 2001. Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation. *J. Med. Genet.* 38: 151-158.
6. Matthews, J.M. and Sunde, M. 2002. Zinc fingers—folds for many occasions. *IUBMB Life* 54: 351-355.
7. Brown, R.S. 2005. Zinc finger proteins: getting a grip on RNA. *Curr. Opin. Struct. Biol.* 15: 94-98.
8. Hall, T.M. 2005. Multiple modes of RNA recognition by zinc finger proteins. *Curr. Opin. Struct. Biol.* 15: 367-373.
9. Gamsjaeger, R., Liew, C.K., Loughlin, F.E., Crossley, M. and Mackay, J.P. 2007. Sticky fingers: zinc-fingers as protein-recognition motifs. *Trends Biochem. Sci.* 32: 63-70.

CHROMOSOMAL LOCATION

Genetic locus: ZNF879 (human) mapping to 5q35.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF879 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF879 shRNA Plasmid (h): sc-155996-SH and ZNF879 shRNA (h) Lentiviral Particles: sc-155996-V as alternate gene silencing products.

For independent verification of ZNF879 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155996A, sc-155996B and sc-155996C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF879 siRNA (h) is recommended for the inhibition of ZNF879 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF879 gene expression knockdown using RT-PCR Primer: ZNF879 (h)-PR: sc-155996-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.