



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


DAB Substrate Buffer

Description:

The DAB Substrate buffer has been developed for applications that require excellent contrast between the chromogen and Hematoxylin counterstain. When in the presence of peroxidase enzyme and DAB chromogen, the substrate helps to produce a brown precipitate that is insoluble in alcohol. The use of liquid components reduces some risks associated with handling powders (ie. dust inhalation), and eliminates waste which often results from using tablets that require a predetermined final volume. This product provides slightly less sensitivity than ScyTek's DAB Substrate (High Sensitivity) catalog#: ACU.

Uses/Limitations:

Not to be taken internally.
For In Vitro Diagnostic use.
Histological applications.
Do not use if reagents become cloudy.
Do not use past expiration date.
Use caution when handling reagents.
Non-Sterile.



Control Tissue:

Any well fixed paraffin embedded or frozen tissue section.

Ordering information regarding individual components on back page!

Availability:

<u>Item #</u>	<u>Component</u>	<u>Volume</u>	<u>Storage</u>
ACC500	DAB Substrate	500 ml	2-8°C
ACC999	DAB Substrate	1000 ml	2-8°C

Bulk Reagent Ordering Information and Current Pricing at www.scytek.com


Precautions:

Avoid contact with skin and eyes.
Harmful if swallowed.
Follow all Federal, State, and local regulations regarding disposal.


Required but not included:

DAB Chromogen (Scytek Catalog# ACB)

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.


Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands

Procedure:

1. Rinse slide in deionized water prior to application of DAB mixture.
2. Combine 50-80µl of DAB Chromogen with 1ml of DAB Substrate and apply to tissue for 5 minutes.

NOTE: Range = 50-80µl of DAB Chromogen / 1 ml of DAB Substrate. 50µl /ml produces optimal staining quality, 80µl/ml produces maximum sensitivity. Combined mixture may be used for up to six hours.


OPTIONAL: For additional sensitivity, add 20µl of DAB Enhancer (cat.# ACM030, not included) to each 1ml of mixed solution.

3. Rinse slide using deionized water.
4. Apply mixture a second time and incubate for another 5 minutes.
5. Rinse slide in two changes of deionized water.
6. Counterstain as desired.
7. Dehydrate through graded alcohol, clear and coverslip in synthetic medium.


References:

1. Kuller, L., Schmidt, A., Mack, H., Durning, M., Birkebak, T., Reiner, M.T., Anderson, D.M., Morton, W.R., Agy, M.B. Systemic and Intestinal Immune Responses to HIV-2₈₇ Infection in Macaca nemestrina. *AIDS Research and Human Retroviruses*. August 2001, Volume 17(12): pages 1191-1204. Doi:10.1089/088922201316912790.
2. Schaefer, K.L., Brachwitz, K., et al. Expression Profiling of t(12;22) Positive Clear Cell Sarcoma of Soft Tissue Cell Lines Reveals Characteristic Up-Regulation of Potential New Marker Genes Including ERBB3. *Cancer Research*, May 15, 2004, Volume 64; page 3395. Doi: 10.1158/0008-5472.CAN-03-0809.
3. Oznurulu, Y., Celik, I., Sur, E., Telatar, T., Ozparlak, H. Comparative Skin Histology of the White New Zealand and Angora Rabbits: Histometrical and Immunohistochemical Evaluations. *Journal of Animal and Veterinary Advances*. 2009, Volume 8(9): pages 1694-1701.
4. Rivas-Arancibia, S., Guevara-Guzman, R., Lopez-Vidal, Y., Rodriguez-Martinez, E., Zanardo-Gomes, M., Angoa-Perez, M., Raisman-Vozari, R. Oxidative Stress Caused by Ozone Exposure Induces Loss of Brain Repair in the Hippocampus of Adult Rats. *Toxicological Sciences*. 2010, Volume 113(1): pages 187-197, doi.1093/toxsci/kfp252.
5. Gupta, P., Dhingra, K.K., Singhal, S., Khurana, N., Mandal, S., Mohta, A. Primary primitive neuroectodermal tumour (PNET) of testis: an unsuspected diagnosis. *Pathology*, February 2010, Volume 42, Issue 2: pages 179-181. Doi: 10.3109/00313020903494029.
6. Lee, J.H., Pyon, J.K., Kim, D.W., Lee, S.H., Nam, H.S., Kim, C.H., Kang, S.G., Lee, Y.J., Park, Y.J., Jeong, D.J., Cho, M.K. Elevated c-Src and c-Yes expression in malignant skin cancers. *Journal of Experimental & Clinical Cancer Research*. 2010, Volume 29: page 116. Doi: 10.1186/1756-9966-29-116.
7. Ozaltin, F., Ibsirlioglu, T., et al. Disruption of PTPRO Causes Childhood-Onset Nephrotic Syndrome. *The American Journal of Human Genetics*. July 15, 2011, Volume 89(1): pages 139-147. Doi: 10.1016/j.ajhg.2011.05.026.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.


Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands