



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Safranin O Solution (For Staining Nuclei)

**Description:** Safranin O has several uses in histology and cytology but is commonly known for counterstaining nuclei red. This formula contains acetic acid providing a lower pH that reduces non-nucleic background and cytoplasmic staining. This solution is not suitable for bacterial (gram) or cartilage staining.


**Results**

Nuclei: Red  
Background: Light Red to Pink

**Uses/Limitations:** Not to be taken internally.  
For In-Vitro Diagnostic use only.  
Histological applications.  
Do not use if reagents become cloudy.  
This protocol has not been tested on and may not be suitable with frozen sections  
Do not use past expiration date.  
Use caution when handling reagents.  
Non-Sterile.

**Control Tissue:** Any well-fixed tissue

**Availability/Contents:**  
Pricing can be found online at [www.scytek.com](http://www.scytek.com)





<u>Item #</u>	<u>Volume</u>
SOH125	125 ml
SOH250	250 ml
SOH500	500 ml
SOH999	1000 ml

*Please contact for OEM and bulk pricing.*

**Precautions:** Avoid contact with skin and eyes.  
Harmful if swallowed.  
Follow all Federal, State, and local regulations regarding disposal.  
Use in chemical fume hood whenever possible.

**Storage:** Room Temperature (18-25°C)

Storage: 18° C  25° C

 ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

CE IVD

EC REP  
Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands

**Notes:**



1. This solution contains acetic acid giving it a moderate acidity and making it unsuitable for bacterial (gram) staining or cartilage staining. It also may not be suitable for other purposes besides staining for nuclei. If user would like to increase staining capability of background and bacteria, the pH will need to be raised.

**Procedure:**

1. Deparaffinize sections if necessary and hydrate to distilled water.
2. Stain tissue with Safranin O solution for 5-10 minutes.
3. Alcohols may remove stain from tissue. Rinse and dehydrate very quickly in absolute ethanol (<30seconds) or alternatively, briefly rinse and airdry slide for greatest retention of stain.

**References:**

1. Charles H. Sawyek (1940) A Standardized Technic for Safranin O, Stain Technology, 15:1, 3-7, DOI: 10.3109/10520294009110324

Storage: 18° C  25° C ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A. Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands