



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Tris Buffered Saline plus Tween 20 (20x) pH 7.4

Description: ScyTek Tris Buffered Saline + Tween 20 (20x Concentrate) pH of 7.4 is an optimal formulation of pH stabilizers, salts and detergents designed to effectively remove excess material from the tissue sample or microtiter plate wells without disrupting the antibody binding reaction. By maintaining the proper buffering environment, unbound components can be washed away without suppressing antigen-antibody binding interactions, thereby reducing nonspecific background and increasing the specific signal. Our Wash buffers do not contain hazardous preservatives such as Azide or Mercury that may interfere with antibody-antigen binding interactions. For your convenience wash buffer is offered in a wide variety of formulations to meet the needs of your specific ELISA application.

Contents: Tris buffered saline in reagent grade water. Tween 20 is added at a diluted concentration of 0.05%, Tris: 0.05M, NaCl: 0.15M. Final pH of diluted buffer is 7.4±0.05.

Availability:	<u>REF #</u>	<u>Volume</u>	<u>Diluted Volume</u>
	TBT500	500 ml	10 Liters
	TBT999	1000 ml	20 Liters
	TBT010	10 Liters	200 Liters
	TBT-20000	20 Liters	400 Liters

Uses/Limitations: Not to be taken internally.
For In-Vitro Diagnostic use.
For professional use.
Histological applications.
Do not use if reagents become cloudy.
Do not use past expiration date.
Use caution when handling reagents.
Non-Sterile.





Ordering Information and Current Pricing at www.scytek.com

Precautions: Avoid contact with skin and eyes.
Harmful if swallowed.
Follow all Federal, State, and local regulations regarding disposal.

Procedure:

1. Pour 50ml of Tris Buffered Saline plus Tween 20 (20x) pH7.4 in mixing flask and add water to final volume of 1000ml.
2. Stir briefly.

Storage: 18° C  25° C

 ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE

EC REP
Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands

References:

1. Heidari M, Wang D, Delekta P, Sun S. Marek's disease virus immunosuppression alters host cellular responses and immune gene expression in the skin of infected chickens. *Veterinary Immunology and Immunopathology*. 2016 Nov 1;180:21-8.
2. Contreras GA, Thelen K, Ayala-Lopez N, Watts SW. The distribution and adipogenic potential of perivascular adipose tissue adipocyte progenitors is dependent on sexual dimorphism and vessel location. *Physiological reports*. 2016 Oct 1;4(19):e12993.
3. Heidari M, Wang D, Fitzgerald SD, Sun S. Severe necrotic dermatitis in the combs of line 63 chickens infected with Marek's disease virus. *Avian Pathology*. 2016 Sep 2;45(5):582-92.
4. France M, Skorich E, Kadrofske M, Swain GM, Galligan JJ. Sex-related differences in small intestinal transit and serotonin dynamics in high-fat-diet-induced obesity in mice. *Experimental physiology*. 2016 Jan 1;101(1):81-99.
5. Atalay F, Ateşoğlu EB, Yıldız S, Firatlı-Tuglular T, Karakuş S, Bayık M. Relationship of P-selectin glycoprotein ligand-1 to prognosis in patients with multiple myeloma. *Clinical Lymphoma Myeloma and Leukemia*. 2015 Mar 31;15(3):164-70.
6. Heidari M, Fitzgerald SD, Zhang H. Immune Responses in Cecal Tonsils of Marek's Disease Virus-Infected Chickens. *Avian diseases*. 2015 Feb 18;59(2):213-26.
7. Weinkopff T, Mackenzie C, Eversole R, Lammie PJ. Filarial excretory-secretory products induce human monocytes to produce lymphangiogenic mediators. *PLoS Negl Trop Dis*. 2014 Jul 10;8(7):e2893.
8. Heidari M, Fitzgerald SD, Zhang H. Marek's disease virus-induced transient cecal tonsil atrophy. *Avian diseases*. 2014 Jan 24;58(2):262-70.
9. Abdul-Ghafar J, Oh SS, Park SM, Wairagu P, Lee SN, Jeong Y, Eom M, Yong SJ, Jung SH. Expression of adiponectin receptor 1 is indicative of favorable prognosis in non-small cell lung carcinoma. *The Tohoku journal of experimental medicine*. 2013;229(2):153-62.
10. Song S, Kole S, Bernier M. A chemical crosslinking method for the analysis of binding partners of heat shock protein-90 in intact cells. *BioTechniques*. 2012 Apr:1.
11. Zipser B, Bello-DeOcampo D, Diestel S, Tai MH, Schmitz B. Mannitou monoclonal antibody uniquely recognizes paucimannose, a marker for human cancer, stemness, and inflammation. *Journal of Carbohydrate Chemistry*. 2012 May 1;31(4-6):504-18.
12. Terrell SP, Origi FC, Agnew D. Glomerulonephropathy in aged captive Key Largo woodrats (*Neotoma floridana smalli*). *Veterinary pathology*. 2012 Jul;49(4):710-6.
13. Liew M, Groll MC, Thompson JE, Call SL, Moser JE, Hoopes JD, Voelkerding K, Wittwer C, Spendlove RS. Validating a custom multiplex ELISA against individual commercial immunoassays using clinical samples. *Biotechniques*. 2007 Mar 1;42(3):327.
14. Bar-On P, Rockenstein E, Adame A, Ho G, Hashimoto M, Masliah E. Effects of the cholesterol-lowering compound methyl-β-cyclodextrin in models of α-synucleinopathy. *Journal of neurochemistry*. 2006 Aug 1;98(4):1032-45.
15. Leppert PC, Kokenyesi R, Klemenich CA, Fisher J. Further evidence of a decorin-collagen interaction in the disruption of cervical collagen fibers during rat gestation. *American journal of obstetrics and gynecology*. 2000 Apr 30;182(4):805-12.

Warranty:

No products or "Instructions For Use (IFU)" are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 18° C



25° C



ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.



EC REP

Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands