

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Instructions For Use TM3-IFU

Rev. Date: 10/29/03

Revision: 2

Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Fax (435) 755-0015 - www.scytek.com

TMB Precipitating Reagent (Standard Sensitivity)

Description: This liquid substrate for peroxidase consists of tetramethylbenzidine (TMB) plus dilute hydrogen

peroxide in a single-reagent stabilized form. The reagent has been specifically formulated for measuring peroxidase in EIA systems. This reagent is stable for long-term storage, and provides

sensitivity equal to, or greater than, that of other commercially available chromogens.

Form: 3,3',5,5'-tetramethylbenzidine

Contents: TMB in a dilute organic solvent, pH 6.0+ 0.5. Hydrogen peroxide is added at a concentration of 0.03%

Storage: Store TMB solution at 2 - 8°C. Storage below this temperature may result in a slight blue tint, which

disappears after returning to the appropriate temperature. Refrigeration is not mandatory. Do not expose reagents to strong light. Keep bottle tightly capped. Reagents are stable for a least 12 months when stored at room temperature, or 18 months when stored at 2 - 8° C. Avoid

contamination of reagents with labware which has not been thoroughly cleaned. A slight yellow tinge may develop over time. This does not affect product performance. Do not use if solution darkens.

Activating Agents: Peroxidase

Light Sensitivity: Negligible for short exposure times

Reaction Time: Approximately 15 minutes (Range 5 - 60 min.)

Reaction pH: Approximately pH 6.0 (Range 5.0 - 7.0)

Reaction Temperature: Room temperature

Precautions: Do not pipet reagents by mouth. Avoid contact with skin and eyes. In case of contact, flush with plenty

water. Dispose of reagents according to local regulations. If disposed of in the sink, flush with plenty of

water.