

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Anti-HSP70 Antibody

Chicken Anti-Human HSP70 Polyclonal Catalog No. SPC-178



Overview

Purification

Product Name
HSP70 Antibody
Description
Chicken Anti-Human HSP70 Polyclonal
Species Reactivity
Human, Mouse, Rat
Applications
WB, ICC/IF, ELISA
Antibody Dilution
WB (1:1000), ICC/IF (1:120); optimal dilutions for assays should be determined by the user.
Host Species
Chicken
Immunogen Species
Human
Immunogen
Full length Human HSP70
Concentration
1 mg/ml
Conjugates
Alkaline Phosphatase, APC, ATTO 390, ATTO 488, ATTO 565, ATTO 594, ATTO 633, ATTO 655, ATTO 680, ATTO 700, Biotin, FITC, HRP, PE/ATTO 594, PerCP, RPE, Streptavidin, Unconjugated
Properties
Storage Buffer
PBS pH7.4, 50% glycerol, 0.09% sodium azide
Storage Temperature
-20°C
Shipping Temperature
Blue Ice or 4°C

PEG Purified
Clonality
Polyclonal
Isotype
lgY
Specificity
Detects ~70kDa.
Cite This Product
Chicken Anti-Human HSP70 Polyclonal (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SPC-178)
Certificate Of Analysis
1 μ g/ml of SMC-178 was sufficient for detection of HSP70 in 20 μ g of heat shocked HeLa cell lysate by colorimetric immunoblot analysis using Goat anti-chicken IgG:HRP as the secondary antibody.
Biological Description
Alternative Names
HSP70 1 Antibody, HSP70 2 Antibody, HSP70.1 Antibody, HSP72 Antibody, HSPA1 Antibody, HSPA1A Antibody, HSPA1B Antibody
Research Areas
Cancer, Heat Shock
Cellular Localization
Cytoplasm
Accession Number
NP_005336.3
NP_005336.3 Gene ID

Scientific Background


P08107

HSP70 genes encode abundant heat-inducible 70-kDa HSPs (HSP70s). In most eukaryotes HSP70 genes exist as part of a multigene family. They are found in most cellular compartments of eukaryotes including nuclei, mitochondria, chloroplasts, the endoplasmic reticulum and the cytosol, as well as in bacteria. The genes show a high degree of conservation, having at least 50% identity (1). The N-terminal two thirds of HSP70s are more conserved than the C-terminal third. HSP70 binds ATP with high affinity and possesses a weak ATPase activity which can be stimulated by binding to unfolded proteins and synthetic peptides (2). When HSC70 (constitutively expressed) present in mammalian cells was truncated, ATP binding activity was found to reside in an N-terminal fragment of 44 kDa which lacked peptide binding capacity. Polypeptide binding ability therefore resided within the C-terminal half (3). The structure of this ATP binding domain displays multiple features of nucleotide binding proteins (4). All HSP70s, regardless of location, bind proteins, particularly unfolded ones. The molecular chaperones of the HSP70 family recognize and bind to nascent polypeptide chains as well as partially folded intermediates of proteins preventing their aggregation and misfolding. The binding of ATP triggers a critical conformational change leading to the release of the bound substrate protein (5). The universal ability of HSP70s to undergo cycles of binding to and release from hydrophobic stretches of partially unfolded proteins determines their role in a great variety of vital intracellular functions such as protein synthesis, protein folding and oligomerization and protein transport. Looking for more information on HSP70? Visit our new HSP70 Scientific Resource Guide at http://www.HSP70.com.


References

- 1. Balashova N. et al. (2005) J Biol Chem 280: 2186-96.
- 2. Boorstein W. R., Ziegelhoffer T. & Craig E. A. (1993) J. Mol. Evol.38 (1): 1-17.
- 3. Rothman J. (1989) Cell 59: 591 -601.
- 4. DeLuca-Flaherty et al. (1990) Cell 62: 875-887.
- 5. Bork P., Sander C. & Valencia A. (1992) Proc. Natl Acad. Sci. USA 89: 7290-7294.
- 6. Fink A.L. (1999) Physiol. Rev. 79: 425-449.


Product Images



Immunocytochemistry/Immunofluorescence analysis using Chicken Anti-Hsp70 Polyclonal Antibody (SPC-178). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Chicken Anti-Hsp70 Polyclonal Antibody (SPC-178) at 1:120 for 12 hours at 4°C. Secondary Antibody: R-PE Goat Anti-Chicken (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Perinuclear region of cytoplasm. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp70 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.



Western blot analysis of Human HeLa cell lysates showing detection of HSP70 protein using Chicken Anti-HSP70 Polyclonal Antibody (SPC-178). Load: 15 μ g protein. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Chicken Anti-HSP70 Polyclonal Antibody (SPC-178) at 1:1000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.



Immunocytochemistry/Immunofluorescence analysis using Chicken Anti-Hsp70 Polyclonal Antibody (SPC-178). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Chicken Anti-Hsp70 Polyclonal Antibody (SPC-178) at 1:120 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Chicken (green) at 1:200 for 2

hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Perinuclear region of cytoplasm. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp70 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.

Product Citations (0)

Currently there are no citations for this product.

Reviews

There are no reviews yet.