



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Maprotiline hydrochloride

Chemical Properties

CAS No. :	10347-81-6
Formula:	C ₂₀ H ₂₄ ClN
Molecular Weight:	313.864
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Maprotiline hydrochloride (Psymion) is a bridged-ring tetracyclic antidepressant that is both mechanistically and functionally similar to the tricyclic antidepressants, including side effects associated with its use.
Targets(IC50)	5-HT Receptor, Adrenergic Receptor, AChR, Norepinephrine, Autophagy, Histamine Receptor
In vitro	Maprotiline inhibits HERG channels expressed in HEK cells with an IC ₅₀ of 5.2 μM and HERG channels expressed in oocytes with an IC ₅₀ of 24 μM. Maprotiline blocks open channels but has no significant effect on closed channels. [1] Maprotiline is a tetracyclic antidepressant, which strongly inhibits the uptake of noradrenaline, though it is notable in its lack of inhibition of serotonergic uptake. Maprotiline also has markedly less pronounced alpha-adrenergic blocking activity than amitriptyline. [2] Maprotiline decreases cell viability in a concentration- and time-dependent manner in Neuro-2a cells. Maprotiline induces apoptosis and increases caspase-3 activation. Maprotiline also induces [Ca(2+)](i) increases which involves the mobilization of intracellular Ca(2+) stored in the endoplasmic reticulum. [3]
In vivo	Maprotiline results in significantly increases GluR1 and GluR2/3 subunit expression in the nucleus accumbens and dorsal striatum of mice as detected by immunohistochemistry; and significantly increases GluR1 and GluR2/3 expression in the hippocampus, as demonstrated by Western blot analysis. [4] Maprotiline has a potent anti-inflammatory effect in rats and this effect is linked to the peripheral and supraspinal actions of the drug. [5] Maprotiline impairs learning when administered before training, but no statistically significant effect is evident when administered after training. [6]

Solubility Information

Solubility	Ethanol: 23 mg/mL (73.3 mM), H ₂ O: < 1 mg/mL (insoluble or slightly soluble), DMSO: 45 mg/mL (143.37 mM), < 1 mg/mL refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.1861 mL	15.9307 mL	31.8613 mL
5 mM	0.6372 mL	3.1861 mL	6.3723 mL
10 mM	0.3186 mL	1.5931 mL	3.1861 mL
50 mM	0.0637 mL	0.3186 mL	0.6372 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Ferrer-Villada T, et al. Eur J Pharmacol,2006, 531(1-3), 1-8.
Arenas MC, et al. Behav Brain Res,2006, 166(1), 150-158.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481