



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Taxifolin

Chemical Properties

CAS No. :	480-18-2
Formula:	C ₁₅ H ₁₂ O ₇
Molecular Weight:	304.25
Appearance:	no data available
Storage:	keep away from direct sunlight
	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Taxifolin (Dihydroquercetin) is a flavonoid in many plants such as <i>Taxus chinensis</i> , Siberian larch, <i>Cedrus deodara</i> and so on.
Targets(IC50)	VEGFR,TNF,Tyrosinase,Adrenergic Receptor,Autophagy
In vivo	Taxifolin is easily metabolized, and its metabolites are prevalent in vivo, although limited data exists on its in vivo metabolism[5]. Metabolites with the same fragment pattern may share the same pharmacophore, exerting the same pharmacological effects as taxifolin through an additive effect on the same drug targets. Thus, taxifolin is bioactive in both its parent form and through its metabolites.

Solubility Information

Solubility	DMSO: 65 mg/mL (213.64 mM), Ethanol: 56 mg/mL (184.1 mM), H ₂ O: 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.2868 mL	16.4339 mL	32.8677 mL
5 mM	0.6574 mL	3.2868 mL	6.5735 mL
10 mM	0.3287 mL	1.6434 mL	3.2868 mL
50 mM	0.0657 mL	0.3287 mL	0.6574 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Abderrezag N, Montenegro Z J S, Louaer O, et al. One-step sustainable extraction of Silymarin compounds of wild Algerian milk thistle (*Silybum marianum*) seeds using Gas Expanded Liquids. Journal of Chromatography A. 2022: