

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# Data Sheet (Cat.No.T2696)



# PRT062607 hydrochloride

Chemical Propert	ties	
CAS No. :	1370261-97-4	
Formula:	C19H23N9Oj¤HCl	
Molecular Weight:	429.91 HG	
Appearance:	no data available	
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year	

## **Biological Description**

Description	PRT062607 hydrochloride (P505-15 Hydrochloride) is a selective inhibitor of Syk (IC50: 1 nM). It displays at least 80-fold selectivity for Syk over other kinases.
Targets(IC50)	MLK,Syk,Src
In vitro	PRT062607 (P505-15) is a highly specific and potent inhibitor of purified Syk (IC50 1-2 nM). In human whole blood, P505-15 potently inhibited B cell antigen receptor- mediated B cell signaling and activation (IC50 0.27 and 0.28 μM, respectively) and Fcε receptor 1-mediated basophil degranulation (IC50 0.15 μM). Similar levels of ex vivo inhibition were measured after dosing in mice (Syk signaling IC50 0.32 μM) [1]. P505-15 abrogated the pro-survival effect of anti-IgM and induced CLL cell apoptosis. Treatment with P505-15 (2 mM/mL) decreased CLL cell viability to 62.9 ± 5.1% after 48 hours. Treatment with lower concentrations of PRT318 and P505-15 also reduced the viability of CLL cells [2].
In vivo	Demonstrating a rapid onset of action, 0.5 h after oral administration of 30 mg/kg P505- 15, Syk activity was reduced by 80% and nearly completely suppressed for the next 5 h relative to vehicle control-treated mice. Syk activity remained more than 60% suppressed for the first 8 h postdosing, returning to levels of vehicle control treatment by 24 h. At the lower dose of 15 mg/kg, more than 50% suppression of Syk activity was observed between 2 and 6 h after oral administration of the compound [1].
Kinase Assay	Potency of Syk inhibition was determined by using a fluorescence resonance energy transfer (FRET) assay. The extent of substrate phosphorylation by Syk was measured in the presence of various P505-15 concentrations. Syk activity was determined by a fluorescent antibody specific for phosphorylated tyrosine by using the increase of FRET. Twelve concentrations were tested for dose response. Specificity and potency of kinase inhibition was determined by evaluation of P505-15 in the Millipore KinaseProfiler panel of 270 independent purified kinase assays. For profiling, P505-15 was tested in duplicate at two concentrations at a fixed concentration of ATP. Subsequently, IC50 determinations using the radioactive assays were carried out at an ATP concentration optimized for each individual kinase. All radioactive ATP incorporation enzyme assays were performed at Millipore UK [1].
Cell Research	NLC co-cultures were established by suspending PBMC from patients with CLL in complete RPMI medium with 10% fetal bovine serum and penicillin-streptomycin-glutamine to a concentration of 107 cells/mL (total 2 mL). Cells were incubated for 14

	days in 24-well plates as previously described. To evaluate whether the Syk inhibitors PRT318 and P505-15 could overcome the protective effect of NLC, CLL cells were cultured under standardized conditions on NLC or in suspension, in the presence or absence of PRT318 and P505-15. At the indicated time points, CLL cells were collected and assayed for cell viability as previously described [2].
Animal Research	All animal studies were performed in strict accordance with the Institutional Animal Care and Use Committee ethical guidelines. Female BALB/c mice received a single oral dose of 15 or 30 mg/kg P505-15 and were anesthetized with a subcutaneous ketamine cocktail, and blood was harvested via cardiac puncture at 0.5, 1, 2, 3, 4, 5, 6, 8, and 24 h postdose, (n = 3/time point; n = 8 vehicle controls). Blood was dispensed into three heparin-containing tubes, one for determination of drug concentration and the remaining two for ex vivo stimulation with isotype control or anti-mouse IgD antibody for 10 min. Blood was processed for intracellular phospho-flow cytometry to evaluate BCR signaling as described earlier; mouse B cells were detected by using CD45R-B220 PerCP-conjugated antibody. Plasma samples were analyzed for P505-15 concentration by using a liquid chromatography tandem mass spectrometer. The analytical range was 2 to 5000 ng/ml [1].

## Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), br/>H2O: 79 mg/mL (183.8 mM),
	 br/>DMSO: 45 mg/mL (104.67 mM), (< 1 mg/ml refers to the product slightly
	soluble or insoluble)

### **Preparing Stock Solutions**

	1mg	5mg	10mg	
1 mM	2.3261 mL	11.6303 mL	23.2607 mL	
5 mM	0.4652 mL	2.3261 mL	4.6521 mL	
10 mM	0.2326 mL	1.163 mL	2.3261 mL	
50 mM	0.0465 mL	0.2326 mL	0.4652 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Coffey G, et al. Specific inhibition of spleen tyrosine kinase suppresses leukocyte immune function and inflammation in animal models of rheumatoid arthritis. J Pharmacol Exp Ther. 2012 Feb;340(2):350-9.

### Inhibitor • Natural Compounds • Compound Libraries • Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481