



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Batabulin sodium

Chemical Properties

CAS No. :	195533-98-3
Formula:	C ₁₃ H ₆ F ₆ NNaO ₃ S
Molecular Weight:	393.24
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Batabulin sodium (T138067 sodium) is an antitumor agent, which covalently and selectively modifies the β -tubulin isotypes, thereby disrupting microtubule polymerization. Batabulin sodium affects cell morphology and leads to cell-cycle arrest ultimately induces apoptotic cell death [1].
In vitro	Treatment with Batabulin (T138067; 30-300 nM; 24-48 hours) in MCF7 cells results in cell-cycle arrest at the G2/M boundary, evidenced by 25-30% of cells displaying tetraploid (4n) DNA content, and induces apoptosis, with 25-30% of cells becoming apoptotic within 24-48 hours, and 50-80% after 48-hour exposure to 100 nM. Batabulin selectively and covalently binds to the β 1, β 2, and β 4 isotypes of β -tubulin at the conserved Cys-239 residue, disrupting microtubule polymerization, leading to cytoskeletal collapse, altered cell shape, and increased chromosomal ploidy. Cell cycle and apoptosis analyses further substantiate these effects, highlighting Batabulin's potential mechanism of action through perturbation of the cell cycle and induction of apoptosis in MCF7 cells [1].
In vivo	Batabulin treatment (T138067; 40 mg/kg; intraperitoneal injection; once per week on days 5, 12, and 19) significantly impaired the growth of drug-sensitive CCRF-CEM tumors in a study involving male athymic nude mice (nu/nu; 6-8 weeks old, 20-25 g) that were injected with CCRF-CEM cells [1]. The study utilized a dosage of 40 mg/kg administered through intraperitoneal injection once per week on specified days, leading to the observed result of hindered tumor growth.

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.543 mL	12.7149 mL	25.4298 mL
5 mM	0.5086 mL	2.543 mL	5.086 mL
10 mM	0.2543 mL	1.2715 mL	2.543 mL
50 mM	0.0509 mL	0.2543 mL	0.5086 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481