



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-SPN Monoclonal Antibody-AF647

Product Details

Ig Type: Rabbit monoclonal IgG
Reactivity: Human
Molecular Weight:
Purification:

Applications

Verified Activity: Flow cytometry analysis of SPN overexpressed 441F cells with TMAZ-0371A6, followed by goat anti-rabbit IgG-ABflo 647 (red line). The isotype control is rabbit IgG (black line).
Application: ELISA, FACS
Recommended: 0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

Properties

Appearance: no data available
Purity: > 95% as determined by SDS-PAGE.
Stability & Storage: For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping: For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

Antigen Details

Immunogen: SPN
Antigen Species: Human
Gene ID: 6693
Uniprot ID: P16150
Synonyms: Galactoglycoprotein;GPL115;Leukocyte sialoglycoprotein;Leukosialin;Sialophorin
Biology Area: Immunology Research

Research Background

Predominant cell surface sialoprotein of leukocytes which regulates multiple T-cell functions, including T-cell activation, proliferation, differentiation, trafficking and migration. Positively regulates T-cell trafficking to lymph-nodes via its association with ERM proteins (EZR, RDX and MSN) (By similarity). Negatively regulates Th2 cell differentiation and predisposes the differentiation of T-cells towards a Th1 lineage commitment. Promotes the expression of IFN-gamma by T-cells during T-cell receptor (TCR) activation of naive cells and induces the expression of IFN-gamma by CD4(+) T-cells and to a lesser extent by CD8(+) T-cells (PubMed:18036228). Plays a role in preparing T-cells for cytokine sensing and differentiation into effector cells by inducing the expression of cytokine receptors IFNGR and IL4R, promoting IFNGR and IL4R signaling and by mediating the clustering of IFNGR with TCR (PubMed:24328034). Acts as a major E-selectin ligand responsible for Th17 cell rolling on activated vasculature and recruitment during inflammation. Mediates Th17 cells, but not Th1 cells, adhesion to E-selectin. Acts as a T-cell

counter-receptor for SIGLEC1 (By similarity)|Protects cells from apoptotic signals, promoting cell survival

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481