



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CD69 Monoclonal Antibody-APC

## Product Details

Ig Type:	Rabbit monoclonal IgG
Reactivity:	Human
Molecular Weight:	150 kDa
Purification:	Protein A Affinity Purified

## Applications

Verified Activity:	Flow cytometry analysis of CD69 overexpressed 454F cells with TMAZ-0384A, followed by goat anti-rabbit IgG-ABflo 647 (red line). The isotype control is rabbit IgG (black line).
Application:	ELISA, FACS
Recommended	0.1-0.2 µg/10E6 cells for FACS; 1 ng/µl for ELISA

## Properties

Appearance:	no data available
Purity:	> 95% as determined by SDS-PAGE.
Stability & Storage:	For unconjugated and biotin antibody, store at -20°C or -80°C for long term. Avoid repeated freeze-thaw cycles. For APC/PE/AF647/AF555/AF488 conjugated antibody, store at 4°C, do not freeze.
Shipping:	For unconjugated and biotin antibody, shipping with dry ice. For APC/PE/AF647/AF555/AF488 conjugated antibody, shipping with blue ice.

## Antigen Details

Immunogen:	CD69
Antigen Species:	Human
Gene ID:	969
Uniprot ID:	Q07108
Synonyms:	Early T-cell activation antigen p60;EA1;Activation inducer molecule;C-type lectin domain family 2 member C;Early activation antigen CD69;GP32/28;MLR-3;BL-AC/P26;Leukocyte surface antigen Leu-23
Biology Area:	Immunology Research

## Research Background

Transmembrane protein expressed mainly on T-cells resident in mucosa that plays an essential role in immune cell homeostasis. Rapidly expressed on the surface of platelets, T-lymphocytes and NK cells upon activation by various stimuli, such as antigen recognition or cytokine signaling, stimulates different signaling pathways in different cell types (PubMed:24752896, PubMed:26296369, PubMed:35930205). Negatively regulates Th17 cell differentiation through its carbohydrate dependent interaction with galectin-1/LGALS1 present on immature dendritic cells (PubMed:24752896). Association of CD69 cytoplasmic tail with the JAK3/STAT5 signaling pathway regulates the transcription of RORgamma/RORC and, consequently, differentiation toward the Th17 lineage (By similarity). Acts also via the S100A8/S100A9 complex present on peripheral blood mononuclear cells to promote the conversion of naive CD4 T-cells into regulatory T-cells (PubMed:26296369). Acts as an oxidized low-density lipoprotein (oxLDL)

receptor in CD4 T-lymphocytes and negatively regulates the inflammatory response by inducing the expression of PDCD1 through the activation of NFAT (PubMed:35930205). Participates in adipose tissue-derived mesenchymal stem cells (ASCs)-mediated protection against *P. aeruginosa* infection. Mechanistically, specifically recognizes *P. aeruginosa* to promote ERK1 activation, followed by granulocyte-macrophage colony-stimulating factor (GM-CSF) and other inflammatory cytokines secretion (PubMed:34841721). In eosinophils, induces IL-10 production through the ERK1/2 pathway (By similarity). Negatively regulates the chemotactic responses of effector lymphocytes and dendritic cells (DCs) to sphingosine 1 phosphate/S1P by acting as a S1PR1 receptor agonist and facilitating the internalization and degradation of the receptor (PubMed:37039481)

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:36 Washington Street,Wellesley Hills,MA 02481