

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Data Sheet (Cat.No.TN6660)



2-Ethylhexyl trans-4-methoxycinnamate

Ch	em	ical	Pro	perti	ies

CAS No. :	83834-59-7	
Formula:	C18H26O3	
Molecular Weight:	290.4	
Appearance:	no data available	CH ₃
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 yea	ar

biological Description	
Description	2-Ethylhexyl trans-4-methoxycinnamate is a sunscreen agent.
Targets(IC50)	Others
In vitro	This study aims to assess the removal of a set of non-polar pollutants in biologically treated wastewater using ozonation, ultraviolet (UV 254 nm low pressure mercury lamp) and visible light (Xe-arc lamp) irradiation as well as visible light photocatalysis using Ce-doped TiO2. The compounds tracked include UV filters, synthetic musks, herbicides, insecticides, antiseptics and polyaromatic hydrocarbons. METHODS AND RESULTS:Raw wastewater and treated samples were analyzed using stir-bar sorptive extraction coupled with comprehensive two-dimensional gas chromatography (SBSE-CG × GC-TOF-MS). Ozone treatment could remove most pollutants with a global efficiency of over 95% for 209 μ M ozone dosage. UV irradiation reduced the total concentration of the sixteen pollutants tested by an average of 63% with high removal of the sunscreen 2-Ethylhexyl trans-4-methoxycinnamate (EHMC), the synthetic musk 7-acetyl-1,1,3,4,4,6-hexamethyltetrahydronaphthalene (tonalide, AHTN) and several herbicides. Visible light Ce-TiO2 photocatalysis reached ~70% overall removal with particularly high efficiency for synthetic musks. In terms of power usage efficiency expressed as nmol kJ(-1), the results showed that ozonation was by far the most efficient process, ten-fold over Xe/Ce-TiO2 visible light photocatalysis, the latter being in turn considerably more efficient than UV irradiation. In all cases the efficiency decreased along the treatments due to the lower reaction rate at lower pollutant concentration. The use of photocatalysis greatly improved the efficiency of visible light irradiation. The collector area per order decreased from 9.14 ± 5.11 m(2) m(-3) order(-1) for visible light irradiation. CONCLUSIONS: Ozonation reduced the toxicity of treated wastewater, while UV irradiation and visible light photocatalysis limited by 20-25% the algal growth due to the accumulation of reaction by-products. Three transformation products were identified and tracked along the treatments[1].

Solubility Information

A DRUG SCREENING EXPERT

DMSO: 60 mg/mL (206.61 mM) (< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg	
1 mM	3.4435 mL	17.2176 mL	34.4353 mL	
5 mM	0.6887 mL	3.4435 mL	6.8871 mL	
10 mM	0.3444 mL	1.7218 mL	3.4435 mL	
50 mM	0.0689 mL	0.3444 mL	0.6887 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Energy efficiency for the removal of non-polar pollutants during ultraviolet irradiation, visible light photocatalysis and ozonation of a wastewater effluent. Water Res. 2013 Oct 1;47(15):5546-56.

Inhibitor • Natural Compounds • Compound Libraries • Recombinant ProteinsThis product is for Research Use Only• Not for Human or Veterinary or Therapeutic UseTel:781-999-4286E_mail:info@targetmol.comAddress:36 Washington Street,Wellesley Hills,MA 02481