



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Solasodine hydrochloride

Chemical Properties


CAS No. : 6106-33-8

Formula: C₂₇H₄₄ClNO₂

Molecular Weight: 450.1

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	Solasodine hydrochloride (90 μ M; 2 days) treatment induced significant budding in P19 cells. This compound strongly stimulated the expression of various neuronal markers, including β III-tubulin, synaptophysin, MAP2, ChAT, and the neural progenitor marker doublecortin. Predominantly, Solasodine hydrochloride directed the differentiation of P19 cells towards neuronal pathways.
In vitro	Solasodine hydrochloride (90 μ M; 2 days) treatment induced significant budding in P19 cells. This compound strongly stimulated the expression of various neuronal markers, including β III-tubulin, synaptophysin, MAP2, ChAT, and the neural progenitor marker doublecortin. Predominantly, Solasodine hydrochloride directed the differentiation of P19 cells towards neuronal pathways.
In vivo	Solasodine hydrochloride, administered as a single intraperitoneal injection at doses ranging from 25-100 mg/kg, significantly prolongs the latency period of the hind limb tonic extensor (HLTE) phase during convulsions induced by picrotoxin. It also significantly enhances thiopental-induced sleep in a dose-dependent manner. Additionally, when administered via intracerebroventricular injection at 375 μ M consistently over two weeks, Solasodine hydrochloride markedly increases the uptake of bromodeoxyuridine in ependymal layer, subventricular zone, and cortical cells, which colocalize with doublecortin immunostaining. Treatment in rats significantly boosts the expression of cholesterol and drug-binding transport proteins in ependymal cells. The compound exhibits anticonvulsant effects and suppresses central nervous system activity. In animal models, Swiss albino mice weighing 18-25 g were used, treated with either picrotoxin or thiopental.

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2217 mL	11.1086 mL	22.2173 mL
5 mM	0.4443 mL	2.2217 mL	4.4435 mL
10 mM	0.2222 mL	1.1109 mL	2.2217 mL
50 mM	0.0444 mL	0.2222 mL	0.4443 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481