



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Macrophage-activating lipopeptide 2

Chemical Properties

CAS No. :	250718-44-6
Formula:	C99H167N19O30S
Molecular Weight:	2135.56
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Macrophage-activating lipopeptide 2 (MALP-2) is a lipopeptide that acts as an agonist for Toll-like receptors TLR-2 and TLR-6. It enhances the phosphorylation of endothelial nitric oxide synthase (eNOS) and the release of nitric oxide (NO) by endothelial cells, thereby improving vasodilation. Additionally, MALP-2 increases the adhesion of leukocytes to the endothelium and enhances perfusion recovery and collateral growth in the hind limbs of hypercholesterolemic Apoe-deficient mice subjected to experimental femoral artery ligation (FAL).
In vitro	Macrophage-activating lipopeptide 2 (1 µg/mL, 24 h) enhances the expression of primary endothelial adhesion molecules and leukocyte integrin receptors, thereby improving leukocyte endothelial adhesion. Additionally, Macrophage-activating lipopeptide 2 (1 µg/mL, 2 h) promotes the phosphorylation of Protein Kinase B and eNOS in endothelial MyEnd cells, facilitating the release of nitric oxide (NO).
In vivo	Macrophage-activating lipopeptide 2 (1 µg dissolved in 125 µL PBS; administered daily; for 10 days; via intravenous injection) enhances perfusion recovery and collateral growth in the hind limbs of hypercholesterolemic Apoe knockout mice subjected to experimental femoral artery ligation (FAL).

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	0.4683 mL	2.3413 mL	4.6826 mL
5 mM	0.0937 mL	0.4683 mL	0.9365 mL
10 mM	0.0468 mL	0.2341 mL	0.4683 mL
50 mM	0.0094 mL	0.0468 mL	0.0937 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481