

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## **VECTASTAIN® ABC-AP Universal Kit**

Alkaline Phosphatase (Horse Anti-Mouse/Rabbit IgG)



Cat. No.	AK-5200
Storage	Store reagents in original bottles at 2–8 °C. Do not freeze.
Description	Instructions for Immunohistochemical staining.

VECTASTAIN ABC-AP Kits are enzymatic, avidin/biotin based amplification kits that produces crisp, highly sensitive, specific staining with low background.

The reagents are supplied in convenient dropper bottles.

### Kit Components

Product Name	Volume
Normal Horse Serum – yellow-labeled bottle	3 ml
Horse Anti-Mouse/Rabbit IgG, Biotinylated – blue-labeled bottle	2 ml
Reagent A (Avidin, ABC-AP) – gray-labeled bottle	2 ml
Reagent B (Biotinylated AP, ABC-AP) – gray-labeled bottle	2 ml

The VECTASTAIN ABC-AP Universal Kit will stain approximately 1000–2000 sections.


#### Preparation of VECTASTAIN Working Solutions

A number of different buffers can be used in the VECTASTAIN ABC-AP system. One of the most common is 10 mM sodium phosphate, pH 7.5, 0.9% saline (PBS). The VECTASTAIN working solutions are prepared as follows:

- Blocking Serum (Normal Serum): add one (1) drop (50 µl) of stock (yellow label) to 5 ml of buffer. The preferred serum for blocking is prepared from the same animal species in which the biotinylated secondary antibody is made.
- Biotinylated Universal Antibody: add two (2) drops (100 µl) of normal blocking serum stock (yellow label) to 5 ml buffer and then add two (2) drop (100 µl) of biotinylated universal secondary antibody stock (blue label).
- VECTASTAIN ABC-AP Reagent: add one (1) drop (50 μl) of Reagent A (gray label) to 5 ml of buffer. Then add one (1) drop (50 μl) of Reagent B (gray label) to the same container, mix immediately, and allow VECTASTAIN Reagent to stand for about 15–30 minutes before use.

### **Staining Procedure**

1. For paraffin sections, deparaffinize and hydrate through xylenes or other clearing agents and graded alcohol series.



Tumor - Double Label: Cyclin A (m), VECTASTAIN ABC-AP Universal Kit, Vector Blue AP Substrate (blue), CD2- (m), VECTASTAIN Elite® ABC Universal Kit, Vector NovaRED™ HRP Substrate (red)

For frozen sections or cell preparations fix with acetone or an appropriate fixative for the antigen under study, if necessary.

Wash for 5 minutes in tap water.

- 2. If antigen unmasking is required, perform this procedure using a Vector® Antigen Unmasking Solution, Citrate-based, pH 6.0 (H-3300) or Tris-based pH 9.0 (H-3301).
- If quenching of endogenous alkaline phosphatase activity is required, incubate the slides in BLOXALL® Blocking Solution for 10 minutes.
- 4. Wash in buffer for 5 minutes.
- 5. Incubate for 20 minutes with diluted normal blocking serum.
- 6. Tip off excess serum from sections.
- 7. Incubate for 30 minutes with primary antibody (mouse or rabbit) diluted in buffer with 2.5% normal serum.
- 8. Wash for 5 minutes in buffer.
- 9. Incubate for 30 minutes with diluted biotinylated secondary antibody.
- 10. Wash for 5 minutes in buffer.
- 11. Incubate for 30 minutes with prepared VECTASTAIN ABC-AP Reagent.
- 12. Wash for 5 minutes in buffer.
- 13. Incubate section for 20–30 minutes in AP substate solution (not included in kit) until desired stain intensity develops.
- 14. Rinse in tap water.
- 15. Counterstain (optional), clear and mount.

Detailed product listings, specifications, protocols and additional information is available on our website: vectorlabs.com