



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ImmPRESS®-AP Horse Anti-Goat IgG Polymer Detection Kit

Alkaline Phosphatase

Cat. No. MP-5405

Storage Store reagents in original bottles at 2-8 °C. Do not freeze.

Description Instructions for Immunohistochemical staining using goat primary antibodies.

The ImmPRESS-AP Polymer Kit is an enzymatic, non-biotin amplification system that produces crisp, highly sensitive, specific staining with low background.

The reagents in the ImmPRESS-AP Polymer Kit are ready-to-use—no mixing or titering is necessary to obtain optimal staining. Dilution of the reagents or changes in suggested incubation time may affect performance. The reagents are supplied in convenient dropper bottles.

A number of different wash buffers can be used with the ImmPRESS reagents. One of the most common is 10 mM sodium phosphate, pH 7.5, 0.9% saline (PBS). 0.1% Tween 20 detergent may be added to the wash buffer and is especially recommended for use with automated stainers.

Kit Components

Product Name	Volume
Normal Horse Serum, 2.5%	15 ml or 50 ml
ImmPRESS-AP Horse Anti-Goat IgG Polymer Reagent	15 ml or 50 ml


The ImmPRESS-AP Polymer (15 ml) Kit will stain approximately 75-50 sections and the ImmPRESS-AP Polymer (50 ml) Kit will stain approximately 250-500 sections.

Staining Procedure

- For paraffin sections, deparaffinize and hydrate tissue sections through xylenes or other clearing agents and graded alcohol series.

For frozen sections or cell preparations fix with acetone or an appropriate fixative for the antigen under study, if necessary.

Wash for 5 minutes in tap water.



Prostate (FFPE): PSA (goat) detected with ImmPRESS-AP Horse Anti-Goat IgG and ImmPACT® Vector Red Substrate (red). No Counterstain.

- If antigen unmasking is required, perform this procedure using a Vector® Antigen Unmasking Solution, Citrate-based, pH 6.0 (H-3300) or Tris-based, pH 9.0 (H-3301).
- If quenching of endogenous peroxidase activity is required, incubate the sections in BLOXALL® Blocking Solution for 10 minutes.
- Wash in buffer for 5 minutes.
- Incubate sections for 20 minutes with Normal Horse Serum, 2.5%.
- Tip off excess serum from sections.
- Incubate with goat primary antibody diluted in appropriate antibody diluent solution, such as diluted normal horse serum or BSA.
- Wash in buffer for 5 minutes.
- Incubate for 30 minutes with ImmPRESS-AP Polymer Reagent.
- Wash for 2 x 5 minutes in buffer.
- Incubate in alkaline phosphatase substrate solution (not included) until desired stain intensity develops.
- Rinse sections in tap water.
- Counterstain (optional), clear and mount.

Detailed product listings, specifications, protocols and additional information are available on our website: vectorlabs.com