



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Streptavidin, AMCA

SA-5008-1

[Product Images](#)



Short Description

AMCA Streptavidin is produced by conjugating streptavidin with a coumarin fluorescent dye, 7-amino-4-methylcoumarin-3-acetic acid. This derivative excites in the ultraviolet (350 nm) and emits in the visible (450 nm) producing an intense blue fluorescence.

Amplification of fluorescent signals can be easily achieved with our biotinylated secondary antibodies followed by our highly purified fluorochrome-labeled streptavidin or avidin. Using a biotin/avidin or biotin/streptavidin detection system results in an additional layer of amplification over a directly conjugated secondary antibody.

Features:

- Recommended for routine immunofluorescence applications
- Highly purified and possesses very low non-specific binding properties
- Extremely high affinity for biotin
- Has a high fluorochrome to protein ratio
- Compared to conventional primary and secondary fluorescent techniques, can provide greater sensitivity and lower background staining

Additional Information

Unit Size	1 mg
Applications	Immunofluorescence, In situ hybridization, Flow Cytometry/Cell Separation
Recommended Usage	For diluting this product, we recommend a HEPES- or bicarbonate-buffered saline solution, approximately pH 8.2. Avoid using RPMI 1640 or other biotin-containing solutions as diluents. Serum also can contain biotin and should not be added to diluents. The recommended concentration range for use is 10-30 µg/ml.
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide
Maximum Excitation	345-355 nm
Maximum Emission	448-454 nm
Recommended Storage	2-8 °C
Concentration	1.0 mg/ml
Conjugate	AMCA
Color of Fluorescence	Blue
Format	Concentrate

