



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CD98 [CP1-5] Standard Size Ab00361-10.0

This chimeric human antibody was made using the variable domain sequences of the original mouse IgG1 format for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1, Kappa

Clone Number: CP1-5

Alternative Name(s) of Target: oncoprotein CD98; large neutral amino acid transporter; LAT1

UniProt Accession Number of Target Protein: P08195 and Q01650

Published Application(s): ELISA

Published Species Reactivity: Human

Immunogen: Epitope peptide (LLHGDFHAFSAG-GGGCCD) linked to KLH.

Specificity: Binds specifically to human CD98.

Application Notes: This monoclonal antibody specifically binds to CD98, a heterodimeric glycoprotein forming the large neutral amino acid transporter (LAT1) which preferentially transports branched-chain and aromatic amino acids.

Antibody First Published in: Itoh et al. 2009 Antibody epitope peptides as potential inducers of IgG antibodies against CD98 oncoprotein. Cancer Science 2009; 100(1):126-131 [PMID:19018757](#)

Note on publication: Describes the design of peptides based on the epitope of a tumour suppressive anti-CD98 antibody, used to generate further IgG1 antibodies.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.