



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CD26 [OX-61] Bulk Size Ab00568-23.0-BT

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG3 format, for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** OX-61

**Alternative Name(s) of Target:** DPP4; DPP 4; DPPIV; DPP IV; Dipeptidyl peptidase 4; Bile canaliculus domain-specific membrane glycoprotein; GP110 glycoprotein; T-cell activation antigen CD26 ;OX61; MRC OX-61; MRC OX61

**UniProt Accession Number of Target Protein:** P14740

**Published Application(s):** ICC, IP, RIA, WB, FC, IF, IHC

**Published Species Reactivity:** Rat

**Immunogen:** OX-61 was prepared by immunizing mice with rat thoracic duct lymphoid cells that were enriched for dendritic cells.

**Specificity:** OX-61 binds specifically to DPP4. Evidence: tissue distribution of OX-61 and DPP4 are identical; equal MWTs; OX-61-mediated reduction in DPP4 activity from solubilized liver homogenates. DPP4 is a glycoprotein receptor which acts as a positive regulator of T cell co-activation.

**Application Notes:** This antibody can be used for denaturing and non-denaturing WB and developed using peroxidase-conjugated rabbit anti-mouse. It can also be used for IHC of cryostat sections from adult rat liver. Staining is also observed on the brush border of the small intestine and colonic epithelium, T cell areas of lymphoid cells but not B cell follicles

**Antibody First Published in:** Paterson et al. Antigens of activated rat T lymphocytes including a molecule of 50,000 Mr detected only on CD4 positive T blasts. Mol Immunol 24:1281 1987 [PMID:2828930](#)

**Note on publication:** Describes the use of Ox-48 among other antibodies to identify new markers for activated T-cells, which might have expression related to the active Vs quiescent state or may be involved in specific T-cell functions.

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without

added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg /ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.