



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ATP6V0C 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000527-T01

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-ATP6V0C full-length

Host: Human

Theoretical MW (kDa): 17.16


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-ATP6V0C antibody (H00000527-B01) by Western Blots.

SDS-PAGE Gel



ATP6V0C transfected lysate.

Western Blot



Lane 1: ATP6V0C transfected lysate (17.16 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [527](#)

GeneBank [NM_001694.2](#)
Accession#:

Protein [NP_001685.1](#)
Accession#:

Gene Name: ATP6V0C

Gene Alias: ATP6C,ATP6L,ATPL,VATL,Vma3

Gene Description: ATPase, H⁺ transporting, lysosomal 16kDa, V0 subunit c

Omim ID: [108745](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is part of the V0 domain. This gene had the previous symbols of ATP6C and ATP6L. [provided by RefSeq]

Other Designations: ATPase, H⁺ transporting, lysosomal (vacuolar proton pump) 16kD,ATPase, H⁺ transporting, lysosomal 16kD, V0 subunit c,ATPase, H⁺ transporting, lysosomal, 16-KD,ATPase, H⁺ transporting, lysosomal, V0 subunit c,H(+)-transporting two-sector ATPase, 16 kDa sub

Gene Pathway

[Epithelial cell signaling in Helicobacter pylori infection](#) [Lysosome](#) [Metabolic pathways](#)
[Oxidative phosphorylation](#) [Vibrio cholerae infection](#)

Related Disease

[Attention Deficit Disorder with Hyperactivity](#) [Autistic Disorder](#) [NARP](#)