



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNFRSF17 monoclonal antibody (M08), clone 3D1

Catalog # : H00000608-M08

規格 : [100 ug]

List All

Specification

Product Description: Mouse monoclonal antibody raised against a partial recombinant TNFRSF17.

Immunogen: TNFRSF17 (AAH58291.1, 5 a.a. ~ 50 a.a) partial recombinant protein with GST tag.


Sequence: AGQCSQNEYFDSLHACIPCQLRCSSNTPPLTCQRYCNASVTNSVK

Host: Mouse

Reactivity: Human

Isotype: IgG1 Kappa

Quality Control Testing: Antibody Reactive Against Recombinant Protein.



Western Blot detection against Immunogen (31.13 kDa) .


Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Datasheet:  [Download](#)

Applications

Western Blot (Recombinant protein)

 [Protocol Download](#)

ELISA

Gene Information

Entrez GeneID: [608](#)

GeneBank Accession#: [BC058291.1](#)

Protein [AAH58291.1](#)

Application Image

Western Blot (Recombinant protein)

ELISA

Accession#:**Gene Name:** TNFRSF17**Gene Alias:** BCM,BCMA,CD269**Gene Description:** tumor necrosis factor receptor superfamily, member 17**Omim ID:** [109545](#)**Gene Ontology:** [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq]

Other Designations: B cell maturation antigen,B-cell maturation factor,OTTHUMP00000160261**Gene Pathway**[Cytokine-cytokine receptor interaction](#)**Related Disease**[Arthritis](#), [Rheumatoid Arthritis](#), [Colitis](#), [Ulcerative Crohn Disease](#), [Genetic Predisposition to Disease](#), [Hematologic Diseases](#), [Kidney Failure](#), [Chronic Multiple Myeloma](#), [Occupational Diseases](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.