



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNFRSF17 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000608-T02

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-TNFRSF17 full-length

Host: Human

Theoretical MW (kDa): 20.1


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-TNFRSF17 antibody ([H00000608-D01P](#)) by Western Blots.

SDS-PAGE Gel



TNFRSF17 transfected lysate.

Western Blot



Lane 1: TNFRSF17 transfected lysate (20.10 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [608](#)

GeneBank [BC058291](#)
Accession#:

Protein [AAH58291.1](#)
Accession#:

Gene Name: TNFRSF17

Gene Alias: BCM,BCMA,CD269

Gene Description: tumor necrosis factor receptor superfamily, member 17

Omim ID: [109545](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq]

Other Designations: B cell maturation antigen,B-cell maturation factor,OTTHUMP00000160261

Gene Pathway

[Cytokine-cytokine receptor interaction](#)

Related Disease

[Arthritis](#), [Rheumatoid Arthritis](#), [Colitis](#), [Ulcerative Crohn Disease](#), [Genetic Predisposition to Disease](#), [Hematologic Diseases](#), [Kidney Failure](#), [Chronic Multiple Myeloma](#), [Occupational Diseases](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.