

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## **CEACAM1 293T Cell Transient Overexpression Lysate(Denatured)**

**Catalog #**: H00000634-T02 規格:[100 uL]

#### List All



Page 1 of 2 2016/5/20

Western Blot

Gene Information

Entrez GeneID: 634

Gene Bank

NM 001024912.1

Accession#:

Protein

Gene

NP 001020083.1

Accession#:

Gene Name: CEACAM1

Gene Alias: BGP,BGP1,BGPI

Description: glycoprotein)

Omim ID: <u>109770</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a member of the carcinoembryonic antigen (CEA)

carcinoembryonic antigen-related cell adhesion molecule 1 (biliary

gene family, which belongs to the immunoglobulin superfamily. Two subgroups of the CEA family, the CEA cell adhesion molecules and the pregnancy-specific glycoproteins, are located within a 1.2 Mb cluster on the long arm of chromosome 19. Eleven pseudogenes of the CEA cell adhesion molecule subgroup are also found in the cluster. The encoded protein was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. Multiple cellular activities have been attributed to the encoded protein, including roles in the differentiation and arrangement of tissue three-dimensional structure, angiogenesis, apoptosis, tumor suppression, metastasis, and the modulation of innate and adaptive immune responses. Multiple transcript variants encoding different isoforms have been reported, but the full-length nature of only two has been determined. [provided by RefSeq

.....

Other CD66a antigen,antigen CD66,biliary glycoprotein adhesion

Designations: molecule, carcino embryonic antigen-related cell adhesion molecule 1

#### **Related Disease**

Body Weight Diabetes Mellitus, Type 2 Genetic Predisposition to Disease Meningococcal Infections Metabolic Syndrome X Osteoporosis

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/20