



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



## CALR Pre-design Chimera RNAi

Catalog # : H00000811-R01

規格 : [ 10 nmol ] [ 20 nmol ]

List All

### Specification

**Product** Homo sapiens calreticulin (CALR), mRNA.

**Description:**

**Reactivity:** Human

**Supplied Product:** DEPC water

**Target Refseq:** NM\_004343

**Target Region:** 3' Untranslated Region (3'UTR)

**Storage Instruction:** Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to ensure product integrity.

**Note:** Position of the Chimera RNAi.



### Application Image

RNAi Knockdown

### Publication Reference

1. dsCheck: highly sensitive off-target search software for double-stranded RNA-mediated RNA interference.  
Naito Y, Yamada T, Matsumiya T, Ui-Tei K, Saigo K, Morishita S. Nucleic Acids Res. 2005 Jul 1;33(Web Server issue):W589-91.
2. Functional dissection of siRNA sequence by systematic DNA substitution: modified siRNA with a DNA seed arm is a powerful tool for mammalian gene silencing with significantly reduced off-target effect.  
Ui-Tei K, Naito Y, Zenno S, Nishi K, Yamato K, Takahashi F, Juni A, Saigo K. Nucleic Acids Res. 2008 Apr;36(7):2136-51. Epub 2008 Feb 11.
3. Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference.  
Ui-Tei K, Naito Y, Takahashi F, Haraguchi T, Ohki-Hamazaki H, Juni A, Ueda R, Saigo K. Nucleic Acids Res. 2004 Feb 9;32(3):936-48. Print 2004.
4. siDirect: highly effective, target-specific siRNA design software for mammalian RNA interference.  
Naito Y, Yamada T, Ui-Tei K, Morishita S, Saigo K. Nucleic Acids Res. 2004 Jul 1;32(Web Server issue):W124-9.

### Applications

#### RNAi Knockdown

#### Gene Information

Entrez GeneID: 811

Gene Name: CALR

**Gene Alias:** CRT,FLJ26680,RO,SSA,cC1qR

**Gene** calreticulin

**Description:**


**Omim ID:** [109091](#)

**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** Calreticulin is a multifunctional protein that acts as a major Ca(2+)-binding (storage) protein in the lumen of the endoplasmic reticulum. It is also found in the nucleus, suggesting that it may have a role in transcription regulation. Calreticulin binds to the synthetic peptide KLGFKKR, which is almost identical to an amino acid sequence in the DNA-binding domain of the superfamily of nuclear receptors. Calreticulin binds to antibodies in certain sera of systemic lupus and Sjogren patients which contain anti-Ro/SSA antibodies, it is highly conserved among species, and it is located in the endoplasmic and sarcoplasmic reticulum where it may bind calcium. The amino terminus of calreticulin interacts with the DNA-binding domain of the glucocorticoid receptor and prevents the receptor from binding to its specific glucocorticoid response element. Calreticulin can inhibit the binding of androgen receptor to its hormone-responsive DNA element and can inhibit androgen receptor and retinoic acid receptor transcriptional activities in vivo, as well as retinoic acid-induced neuronal differentiation. Thus, calreticulin can act as an important modulator of the regulation of gene transcription by nuclear hormone receptors. Systemic lupus erythematosus is associated with increased autoantibody titers against calreticulin but calreticulin is not a Ro/SS-A antigen. Earlier papers referred to calreticulin as an Ro/SS-A antigen but this was later disproven. Increased autoantibody titer against human calreticulin is found in infants with complete congenital heart block of both the IgG and IgM classes. [provided by RefSeq]

**Other Designations:** Sicca syndrome antigen A (autoantigen Ro; calreticulin),autoantigen Ro

### Interactome



### Gene Pathway

[Antigen processing and presentation](#)

### Related Disease

[Cardiovascular Diseases](#) [Diabetes Mellitus, Type 2](#) [Edema](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.