



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RUNX1T1 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000862-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-RUNX1T1 full-length

Host: Human

Theoretical MW (kDa): 63.58


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-RUNX1T1 antibody (H00000862-B02) by Western Blots.

SDS-PAGE Gel



RUNX1T1 transfected lysate.

Western Blot



Lane 1: RUNX1T1 transfected lysate (63.58 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [862](#)

GeneBank Accession#: [NM_004349.2](#)

Protein Accession#: =

Gene Name: RUNX1T1

Gene Alias: AML1T1,CBFA2T1,CDR,ETO,MGC2796,MTG8,MTG8b,ZMYND2

Gene Description: runt-related transcription factor 1; translocated to, 1 (cyclin D-related)

Omim ID: [133435](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a putative zinc finger transcription factor and oncoprotein. In acute myeloid leukemia, especially in the M2 subtype, the t(8;21)(q22;q22) translocation is one of the most frequent karyotypic abnormalities. The translocation produces a chimeric gene made up of the 5'-region of the RUNX1 gene fused to the 3'-region of this gene. The chimeric protein is thought to associate with the nuclear corepressor/histone deacetylase complex to block hematopoietic differentiation. Several transcript variants encoding multiple isoforms have been found for this gene. [provided by RefSeq]

Other Designations: acute myelogenous leukemia 1 translocation 1 protein,acute myelogenous leukemia 1 translocation 1, cyclin-D related,core-binding factor, runt domain, alpha subunit 2; translocated to, 1; cyclin D-related,eight twenty one protein,myeloid translocation gene

Gene Pathway

[Acute myeloid leukemia Pathways in cancer](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.