



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com



[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CCNA2 polyclonal antibody (A01)

Catalog # : H00000890-A01

規格 : [50 uL]

List All

Specification	Application Image
Product Description: Mouse polyclonal antibody raised against a partial recombinant CCNA2.	ELISA
Immunogen: CCNA2 (NP_001228, 1 a.a. ~ 100 a.a) partial recombinant protein with GST tag.	
Sequence: MLGNSAPGPATREAGSALLALQQTALQEDQENINPEKAAPVQQPRTRA ALAVLKSGNPRGLAQQQRPKTRRRVAPLKDLPVNDEHVTVPPWKANSKQ PAFT	
Host: Mouse	
Reactivity: Human	
Quality Control Testing: Antibody Reactive Against Recombinant Protein.	
Storage Buffer: 50 % glycerol	
Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.	
MSDS:  Download	
Datasheet:  Download	
Applications	
ELISA	
Gene Information	
Entrez GeneID: 890	
GeneBank Accession#: NM_001237	
Protein Accession#: NP_001228	
Gene Name: CCNA2	
Gene Alias: CCN1,CCNA	
Gene Description: cyclin A2	
Omim ID: 123835	
Gene Ontology: Hyperlink	

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. In contrast to cyclin A1, which is present only in germ cells, this cyclin is expressed in all tissues tested. This cyclin binds and activates CDC2 or CDK2 kinases, and thus promotes both cell cycle G1/S and G2/M transitions. [provided by RefSeq]

Other cyclin A

Designations:

Gene Pathway

[Cell cycle](#)

Related Disease

[Adenocarcinoma Esophageal Neoplasms Genetic Predisposition to Disease](#)
[Kidney Failure, Chronic Lung Neoplasms Ovarian Neoplasms](#)
[Pulmonary Disease, Chronic Obstructive Urinary Bladder Neoplasms Werner syndrome](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.