

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





# **CCNE1** polyclonal antibody (A01)

**Catalog #**: H00000898-A01 規格:[50 uL]

### List All

Product Description:	Mouse polyclonal antibody raised against a partial recombinant CCNE1
lmmunogen:	CCNE1 (NP_001229, 311 a.a. ~ 410 a.a) partial recombinant protein with GST tag.
Sequence:	ELMQKVSGYQWCDIENCVKWMVPFAMVIRETGSSKLKHFRGVADEDAH NIQTHRDSLDLLDKARAKKAMLSEQNRASPLPSGLLTPPQSGKKQSSGP EMA
Host:	Mouse
Reactivity:	Human
Quality Control Testing:	Antibody Reactive Against Recombinant Protein.  175- 83- 62- 47.5- 32- Western Blot detection against Immunogen (37.11 KDa).
Storage Buffer:	50 % glycerol
Storage Instruction:	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
MSDS:	Download
Datasheet:	Download
expression pr Hudelist G, Si	alysis in human breast cancer: identification of a characteristic protein rofile of malignant breast epithelium. inger CF, Pischinger KI, Kaserer K, Manavi M, Kubista E, Czerwenka ss. 2006 Mar;6(6):1989-2002.
Applications	

### **Application Image**

Western Blot (Recombinant protein)

ELISA

**ELISA** 

**Gene Information** 

Entrez GeneID: 898

**GeneBank** 

NM\_001238

Accession#:

Protein

NP\_001229

Accession#:

Gene Name: CCNE1

Gene Alias: **CCNE** 

Gene

cyclin E1

**Description:** 

Omim ID: 123837


Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. Two alternatively spliced transcript variants of this gene, which encode distinct isoforms, have been described. Two additional splice variants were reported but detailed nucleotide sequence information is not yet available. [provided by RefSeq

Other **Designations:**  cyclin Es,cyclin Et

Interactome

Page 2 of 3 2016/5/20



#### **Gene Pathway**

<u>Cell cycle p53 signaling pathway Pathways in cancer Prostate cancer</u> Small cell lung cancer

#### **Related Disease**

Adenocarcinoma Breast cancer Breast Neoplasms Disease Progression
Esophageal Neoplasms Genetic Predisposition to Disease Neoplasm Invasiveness
Neoplasms, Glandular and Epithelial Ovarian cancer Ovarian Neoplasms
Urinary Bladder Neoplasms

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20