

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



CCNE1 MaxPab mouse polyclonal antibody (B01) MaxPab®

規格:[50 uL]

Catalog # : H00000898-B01

List All

Specification		Application Image
Product Description:	Mouse polyclonal antibody raised against a full-length human CCNE1 protein.	Western Blot (Transfected lysate)
Immunogen:	CCNE1 (AAH35498, 1 a.a. ~ 410 a.a) full-length human protein.	100 - 75 - 50 - 37 -
Sequence:	MPRERRERDAKERDTMKEDGGAEFSARSRKRKANVTVFLQDPDEETA KIDRTARDQCGSQPWDNNAVCADPCSLIPTPDKEDDDRVYPNSTCKPRII APSRGSPLPVLSWANREEVWKIMLNKEKTYLRDQHFLEQHPLLQPKMRA ILLDWLMEVCEVYKLHRETFYLAQDFFDRYMATQENVVKTLLQLIGISSLF IAAKLEEIYPPKLHQFAYVTDGACSGDEILTMELMIMKALKWRLSPLTIVS WLNVYMQVAYLNDLHEVLLPQYPQQIFIQIAELLDLCVLDVDCLEFPYGIL AASALYHFSSSELMQKVSGYQWCDIENCVKWMVPFAMVIRETGSSKLKH FRGVADEDAHNIQTHRDSLDLLDKARAKKAMLSEQNRASPLPSGLLTPP QSGKKQSSGPEMA	25- 20- 15- enlarge
Host:	Mouse	
Reactivity:	Human	
Quality Control Testing:	Antibody reactive against mammalian transfected lysate.	
Storage Buffer	: No additive	
Storage Instruction:	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.	
Note:	For IHC and IF applications, antibody purification with Protein A will be needed prior to use.	
MSDS:	Download	

Western Blot (Transfected lysate)



Western Blot analysis of CCNE1 expression in transfected 293T cell line (<u>H00000898-T01</u>) by CCNE1 MaxPab polyclonal antibody.

Gene Information		
Entrez GenelD: <u>898</u>		
GeneBank Accession#:		
Protein Accession#:	<u>AAH35498</u>	
Gene Name:		
	CCNE	
Gene Description:	cyclin E1	
Omim ID:	<u>123837</u>	

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. Two alternatively spliced transcript variants of this gene, which encode distinct isoforms, have been described. Two additional splice variants were reported but detailed nucleotide sequence information is not yet available. [provided by RefSeq

Other cyclin Es,cyclin Et

Designations:

Interactome



Gene Pathway

<u>Cell cycle p53 signaling pathway Pathways in cancer</u> <u>Prostate cancer</u> <u>Small cell lung cancer</u>

Related Disease

Adenocarcinoma Breast cancer Breast Neoplasms Disease Progression Esophageal Neoplasms Genetic Predisposition to Disease Neoplasm Invasiveness Neoplasms, Glandular and Epithelial Ovarian cancer Ovarian Neoplasms Urinary Bladder Neoplasms

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.