

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





CCNE1 Pre-design Chimera RNAi

List All

cification								Application Image RNAi Knockdown
Homo sapiens cyclin E1 (CCNE1), transcript variant 2, mRNA.								
Human								
DEPC wat	ter							
: NM_0571	82							
Storage Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to ensure product integrity.								
Note: Position of the Chimera RNAi.								
sapiens)								
364 546	727	909	1091	1272	1454	1636	1818	
	Human DEPC wat : NM_0571 Store at -2 product in Position of sapiens)	Human DEPC water : NM_057182 Store at -20°C, do product integrity. Position of the Chinesapiens)	Human DEPC water : NM_057182 Store at -20°C, do not exc product integrity. Position of the Chimera RI sapiens)	Human DEPC water : NM_057182 Store at -20°C, do not exceed 4 - 5 product integrity. Position of the Chimera RNAi.	Human DEPC water : NM_057182 Store at -20°C, do not exceed 4 - 5 freeze product integrity. Position of the Chimera RNAi.	Human DEPC water : NM_057182 Store at -20°C, do not exceed 4 - 5 freeze-thaw cyproduct integrity. Position of the Chimera RNAi. sapiens)	Human DEPC water : NM_057182 Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to product integrity. Position of the Chimera RNAi. sapiens)	Human DEPC water : NM_057182 Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to ensure product integrity. Position of the Chimera RNAi. sapiens)

- dsCheck: highly sensitive off-target search software for double-stranded RNA-mediated RNA interference.
 - Naito Y, Yamada T, Matsumiya T, Ui-Tei K, Saigo K, Morishita S.Nucleic Acids Res. 2005 Jul 1;33(Web Server issue):W589-91.
- Functional dissection of siRNA sequence by systematic DNA substitution: modified siRNA with a DNA seed arm is a powerful tool for mammalian gene silencing with significantly reduced off-target effect.
 - Ui-Tei K, Naito Y, Zenno S, Nishi K, Yamato K, Takahashi F, Juni A, Saigo K.Nucleic Acids Res. 2008 Apr;36(7):2136-51. Epub 2008 Feb 11.
- 3. <u>Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference.</u>
 - Ui-Tei K, Naito Y, Takahashi F, Haraguchi T, Ohki-Hamazaki H, Juni A, Ueda R, Saigo K.Nucleic Acids Res. 2004 Feb 9;32(3):936-48. Print 2004.
- 4. siDirect: highly effective, target-specific siRNA design software for mammalian RNA interference.
 - Naito Y, Yamada T, Ui-Tei K, Morishita S, Saigo K.Nucleic Acids Res. 2004 Jul 1;32(Web Server issue):W124-9.

Applications

RNAi Knockdown

Gene Information

Entrez GenelD: 898

Gene Name: CCNE1

Gene Alias: CCNE

Page 1 of 3 2016/5/20

Gene cyclin E1

Description:


Omim ID: 123837

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. Two alternatively spliced transcript variants of this gene, which encode distinct isoforms, have been described. Two additional splice variants were reported but detailed nucleotide sequence information is not yet available. [provided by

Other **Designations:** cyclin Es, cyclin Et

Interactome



Gene Pathway

Cell cycle p53 signaling pathway Pathways in cancer Prostate cancer Small cell lung cancer

Related Disease

Adenocarcinoma Breast cancer Breast Neoplasms Disease Progression Esophageal Neoplasms Genetic Predisposition to Disease Neoplasm Invasiveness Neoplasms, Glandular and Epithelial Ovarian cancer Ovarian Neoplasms Urinary Bladder Neoplasms

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20