



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CCNH 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000902-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-CCNH full-length

Host: Human

Theoretical MW (kDa): 37.6


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-CCNH antibody (H00000902-B01) by Western Blots.

SDS-PAGE Gel



CCNH transfected lysate

Western Blot



Lane 1: CCNH transfected lysate (37.6 KDa).

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [902](#)

GeneBank Accession#: [NM_001239](#)

Protein Accession#: [NP_001230](#)

Gene Name: CCNH

Gene Alias: CAK,p34,p37

Gene Description: cyclin H

Omim ID: [601953](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. [provided by RefSeq]

Other Designations: CDK-activating kinase,MO15-associated protein,cyclin-dependent kinase-activating kinase

Gene Pathway

[Cell cycle](#) [Nucleotide excision repair](#)

Related Disease

[Adenoma](#) [Ataxia telangiectasia](#) [Biliary Tract Neoplasms](#) [Brain Neoplasms](#)
[Carcinoma, Non-Small-Cell Lung](#) [Colorectal Neoplasms](#) [Esophageal Neoplasms](#)
[Genetic Predisposition to Disease](#) [Glioma](#) [Kidney Failure, Chronic](#)
[Leukemia, Myeloid, Acute](#) [Lung Neoplasms](#) [Meningioma](#) [Mouth Neoplasms](#)
[Multiple Sclerosis](#) [Neoplasm Invasiveness](#) [Neoplasms](#) [Neuroma, Acoustic](#) [Ovarian cancer](#)

... see more