



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MS4A3 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000932-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-MS4A3 full-length

Host: Human

Theoretical MW (kDa): 23.65


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-MS4A3 antibody (H00000932-B01) by Western Blots.

SDS-PAGE Gel



MS4A3 transfected lysate

Western Blot



Lane 1: MS4A3 transfected lysate (23.65 KDa).

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [932](#)

GeneBank Accession#: [BC008487](#)

Protein Accession#: [AAH08487](#)

Gene Name: MS4A3

Gene Alias: CD20L,HTM4

Gene Description: membrane-spanning 4-domains, subfamily A, member 3 (hematopoietic cell-specific)

Omim ID: [606498](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a member of the membrane-spanning 4A gene family. Members of this protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This family member likely plays a role in signal transduction and may function as a subunit associated with receptor complexes. The gene encoding this protein is localized to 11q12, among a cluster of related family members. Alternative splicing may result in multiple transcript variants; however, not all variants have been fully described. [provided by RefSeq]

Other Designations: CD20 antigen homolog,CD20 antigen-like,IgE receptor beta chain,IgE receptor beta subunit,hematopoietic cell 4 transmembrane protein,membrane-spanning 4-domains subfamily A member 3,membrane-spanning 4-domains, subfamily A, member 3

Related Disease

[Genetic Predisposition to Disease](#) [Narcolepsy](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.