



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

TNFRSF7 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00000939-T02

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-TNFRSF7 full-length

Host: Human

Theoretical MW (kDa): 29.2


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-TNFRSF7 antibody ([H00000939-D01P](#)) by Western Blots.

SDS-PAGE Gel



CD27 transfected lysate.

Western Blot



Lane 1: CD27 transfected lysate (29.20 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [939](#)

GeneBank Accession#: [NM_001242](#)

Protein Accession#: [NP_001233.1](#)

Gene Name: CD27

Gene Alias: MGC20393,S152,T14,TNFRSF7,Tp55

Gene Description: CD27 molecule

Omim ID: [186711](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor. [provided by RefSeq]

Other Designations: CD27 antigen,CD27L receptor,T cell activation antigen CD27,T cell antivation antigen S152,tumor necrosis factor receptor superfamily, member 7

Gene Pathway

[Cytokine-cytokine receptor interaction](#)

Related Disease

[Asthma](#) [Asthma](#) [Bronchial Hyperreactivity](#) [Genetic Predisposition to Disease](#)
[Hematologic Diseases](#) [Kidney Failure](#), [Chronic Multiple Myeloma](#) [Occupational Diseases](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.