



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CTSL (Human) Recombinant Protein (P01)

Catalog Number: H00001514-P01

Regulation Status: For research use only (RUO)

Product Description: Human CTSL full-length ORF (AAH12612.1, 1 a.a. - 333 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MNPTLILAAFCGLIASATLTFDHSLEAQWTKWKAMHNR
LYGMNEEGWRRRAVWEKNVKMIELHNQEYREGKHSFT
MAMNAFGDMTSEEFQVMNGFQNRKPRKGKVFQEP
LFYEAPRSVDWREKGYVTPVKNQGGCGSCWAFSAT
GALEGQMFRTGRLISLSEQNLVDCSGPQGNEGCNG
GLMDYAFQYVQDNGGLDSEESYPYEATEESCKYNPK
YSVANDTGFVDIPKQEKALMKAVATVGPISVAIDAGHE
SFLFYKEGIYFEPDCSSEDMDHGVLVVGYGFFESTESD
NNKYWLVKNSWGEEWGMGGYVKMAKDRRNHCGIAS
AASYPTV

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 62.37

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1514

Gene Symbol: CTSL1

Gene Alias: CATL, CTSL, FLJ31037, MEP

Gene Summary: The protein encoded by this gene is a lysosomal cysteine proteinase that plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. The encoded protein has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. This protein, which is a member of the peptidase C1 family, is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. At least two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq]