



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DAB2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00001601-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-DAB2 full-length

Host: Human

Theoretical MW (kDa): 84.81


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-DAB2 antibody (H00001601-B01) by Western Blots.

SDS-PAGE Gel



DAB2 transfected lysate.

Western Blot



Lane 1: DAB2 transfected lysate (84.81 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [1601](#)

GeneBank [BC003064.2](#)
Accession#:

Protein [AAH03064.1](#)
Accession#:

Gene Name: DAB2

Gene Alias: DOC-2,DOC2,FLJ26626

Gene Description: disabled homolog 2, mitogen-responsive phosphoprotein (Drosophila)

Omim ID: [601236](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: DAB2 mRNA is expressed in normal ovarian epithelial cells but is down-regulated or absent from ovarian carcinoma cell lines. The 770-amino acid predicted protein has an overall 83% identity with the mouse p96 protein, a putative mitogen-responsive phosphoprotein; homology is strongest in the amino-terminal end of the protein in a region corresponding to the phosphotyrosine interaction domain. The down-regulation of DAB2 may play an important role in ovarian carcinogenesis. This gene was initially named DOC2 (for Differentially expressed in Ovarian Cancer) and is distinct from the DOC2A and DOC2B genes (for double C2-like domains, alpha and beta). [provided by RefSeq]

Other Designations: disabled homolog 2,mitogen-responsive phosphoprotein

Gene Pathway

[Endocytosis](#)

Related Disease

[Alzheimer Disease](#) [Alzheimer disease](#) [Birth Weight](#) [Cerebral Amyloid Angiopathy](#) [Genetic Predisposition to Disease](#) [Glioblastoma](#) [Glioma](#) [Kidney Failure](#) [Chronic Leukemia](#) [Leukemia](#) [Myeloid, Acute](#) [Meningeal Neoplasms](#) [Meningioma](#) [Neuroblastoma](#) [Ovarian Neoplasms](#) [Tobacco Use Disorder](#)