



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

DCK MaxPab rabbit polyclonal antibody (D03)

Catalog Number: H00001633-D03

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against a full-length human DCK protein.

Immunogen: DCK (NP_000779.1, 1 a.a. ~ 260 a.a) full-length human protein.

Sequence:

```
MATPPKRSCPSFSASSEGTRIKKISIEGNIAAGKSTFVN  
ILKQLCEDWEVVPEPVARWCNVQSTQDEFEELTMSQ  
KNGGNVLQMMYEKPERWSFTFQTYACLSRIRAQLASL  
NGKCLKDAEKPVLFERSVYSDRYIFASNLYESECMNET  
EWTIYQDWHDMNMQFGQSLDGIYQLQATPETCLH  
RIYLRGRNEEQGIPLEYLEKLHYKHESWLLHRTLKTNF  
DYLQEVPIITLDVNEDFKDKYESLVEKVKEFLSTL
```

Host: Rabbit

Reactivity: Human

Applications: IP, WB-Ce, WB-Ti, WB-Tr
(See our web site product page for detailed applications information)

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Buffer: No additive

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1633

Gene Symbol: DCK

Gene Alias: MGC117410, MGC138632

Gene Summary: Deoxycytidine kinase (DCK) is required for the phosphorylation of several deoxyribonucleosides and their nucleoside analogs. Deficiency of DCK is associated with resistance to antiviral and anticancer chemotherapeutic agents.

Conversely, increased deoxycytidine kinase activity is associated with increased activation of these compounds to cytotoxic nucleoside triphosphate derivatives. DCK is clinically important because of its relationship to drug resistance and sensitivity. [provided by RefSeq]