



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DFFA 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00001676-T02

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-DFFA full-length

Host: Human

Theoretical MW (kDa): 36.5


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-DFFA antibody (H00001676-D01) by Western Blots.

SDS-PAGE Gel



DFFA transfected lysate.

Western Blot



Lane 1: DFFA transfected lysate (36.5 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [1676](#)

GeneBank Accession#: [NM_004401](#)

Protein Accession#: [NP_004392.1](#)

Gene Name: DFFA

Gene Alias: DFF-45,DFF1,ICAD

Gene Description: DNA fragmentation factor, 45kDa, alpha polypeptide


Omim ID: [601882](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq]

Other Designations: DFF45,DNA fragmentation factor, 45 kD, alpha polypeptide,DNA fragmentation factor, 45 kD, alpha subunit,OTTHUMP00000001903,OTTHUMP00000001904

Interactome



Gene Pathway

[Apoptosis](#)

Related Disease

[Celiac Disease](#) [Genetic Predisposition to Disease](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.