



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DFFB Pre-design Chimera RNAi

Catalog # : H00001677-R02

規格 : [10 nmol] [20 nmol]

List All

Specification

Product Description: Homo sapiens DNA fragmentation factor, 40kDa, beta polypeptide (caspase-activated DNase) (DFFB), transcript variant 2, mRNA.

Reactivity: Human

Supplied Product: DEPC water

Target Refseq: NM_001004286

Target Region: 3' Untranslated Region (3'UTR)

Storage Instruction: Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to ensure product integrity.

Note: Position of the Chimera RNAi.
The related RNAi products listed below were designed from different accession number but sharing the same RNAi sequence.



Application Image

RNAi Knockdown

Publication Reference

- dsCheck: highly sensitive off-target search software for double-stranded RNA-mediated RNA interference.
Naito Y, Yamada T, Matsumiya T, Ui-Tei K, Saigo K, Morishita S. *Nucleic Acids Res.* 2005 Jul 1;33(Web Server issue):W589-91.
- Functional dissection of siRNA sequence by systematic DNA substitution: modified siRNA with a DNA seed arm is a powerful tool for mammalian gene silencing with significantly reduced off-target effect.
Ui-Tei K, Naito Y, Zenno S, Nishi K, Yamato K, Takahashi F, Juni A, Saigo K. *Nucleic Acids Res.* 2008 Apr;36(7):2136-51. Epub 2008 Feb 11.
- Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference.
Ui-Tei K, Naito Y, Takahashi F, Haraguchi T, Ohki-Hamazaki H, Juni A, Ueda R, Saigo K. *Nucleic Acids Res.* 2004 Feb 9;32(3):936-48. Print 2004.
- siDirect: highly effective, target-specific siRNA design software for mammalian RNA interference.
Naito Y, Yamada T, Ui-Tei K, Morishita S, Saigo K. *Nucleic Acids Res.* 2004 Jul 1;32(Web Server issue):W124-9.

Applications

RNAi Knockdown

Gene Information

Entrez GeneID: [1677](#)

Gene Name: DFFB

Gene Alias: CAD,CPAN,DFF-40,DFF2,DFF40

Gene Description: DNA fragmentation factor, 40kDa, beta polypeptide (caspase-activated DNase)

Omim ID: [601883](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene but the biological validity of these variants has not been determined. [provided by RefSeq]

Other Designations: DNA fragmentation factor, 40 kD, beta polypeptide,DNA fragmentation factor, 40 kD, beta polypeptide (caspase-activated DNase),DNA fragmentation factor, 40 kD, beta subunit,OTTHUMP00000003633,caspase-activated deoxyribonuclease,caspase-activated nuclease

Gene Pathway

[Apoptosis](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.