



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

DR1 (Human) Recombinant Protein (P01)

Catalog Number: H00001810-P01

Regulation Status: For research use only (RUO)

Product Description: Human DR1 full-length ORF (AAH02809, 1 a.a. - 176 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MASSSGNDDDLTIPRAAINKMIKETLPNVRVANDAREL
VVNCCTEFIHLLISSEANEICNKSEKKTISPEHVIALES
LGFSGYISEVKEVLQECKTVALKRRKASSRLENLGIPEE
ELLRQQQELFAKARQQQAELAQQEWLQMQQAAQQA
QLAAASASASNQAGSSQDEEDDDDI

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 45.10

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1810

Gene Symbol: DR1

Gene Alias: NC2, NC2-BETA

Gene Summary: This gene encodes a TBP- (TATA box-binding protein) associated phosphoprotein that represses both basal and activated levels of

transcription. The encoded protein is phosphorylated in vivo and this phosphorylation affects its interaction with TBP. This protein contains a histone fold motif at the amino terminus, a TBP-binding domain, and a glutamine- and alanine-rich region. The binding of DR1 repressor complexes to TBP-promoter complexes may establish a mechanism in which an altered DNA conformation, together with the formation of higher order complexes, inhibits the assembly of the preinitiation complex and controls the rate of RNA polymerase II transcription. [provided by RefSeq]