



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



E2F4 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00001874-T02 規格 : [100 uL]

[List All](#)

Specification

Transfected 293T

Cell Line:

Plasmid: pCMV-E2F4 full-length


Host: Human

Theoretical MW 44


(kDa):

Quality Control Testing: Transient overexpression cell lysate was tested with Anti-E2F4 antibody (H00001874-D01P) by Western Blots.

SDS-PAGE Gel



Western Blot



Lane 1: E2F4 transfected lysate (44.00 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [1874](#)

GeneBank [NM_001950.3](#)

Accession#:

Protein [NP_001941.2](#)

Accession#:

Gene Name: E2F4

Gene Alias: E2F-4

Gene E2F transcription factor 4, p107/p130-binding

Description:

Omim ID: [600659](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionarily conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein binds to all three of the tumor suppressor proteins pRB, p107 and p130, but with higher affinity to the last two. It plays an important role in the suppression of proliferation-associated genes, and its gene mutation and increased expression may be associated with human cancer. [provided by RefSeq]

Other E2F transcription factor 4,p107/p130-binding protein

Designations:

Gene Pathway

[Cell cycle](#) [TGF-beta signaling pathway](#)

Related Disease

[Genetic Predisposition to Disease](#) [Ovarian Neoplasms](#)