



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

EBI2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00001880-T01

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-EBI2 full-length

Host: Human

Theoretical MW (kDa): 39.82


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-EBI2 antibody (H00001880-B01) by Western Blots.

SDS-PAGE Gel



EBI2 transfected lysate

Western Blot



Lane 1: EBI2 transfected lysate (39.82 KDa).

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [1880](#)

**GeneBank
Accession#:** [BC020752](#)

**Protein
Accession#:** [AAH20752](#)

Gene Name: GPR183

Gene Alias: EBI2

**Gene
Description:** G protein-coupled receptor 183

Omim ID: [605741](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene was identified by the up-regulation of its expression upon Epstein-Barr virus infection of primary B lymphocytes. This gene is predicted to encode a G protein-coupled receptor that is most closely related to the thrombin receptor. Expression of this gene was detected in B-lymphocyte cell lines and lymphoid tissues but not in T-lymphocyte cell lines or peripheral blood T lymphocytes. The function of this gene is unknown. [provided by RefSeq]

**Other
Designations:** EBV-induced G protein-coupled receptor 2, Epstein-Barr virus induced gene 2 (lymphocyte-specific G protein-coupled receptor), OTTHUMP00000018611

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.